

Cosmic inflation, deceleration, acceleration, dark matter, and dark ‘energy’ in one coherent package

Homer G. Ellis

*Department of Mathematics, University of Colorado at Boulder, Boulder, Colorado 80309**

(Dated: January 24, 2010)

In creating his gravitational field equations Einstein assumed without justification that inertial mass, even in its equivalent form as energy, is a source of gravity. Denying that assumption allows modifying the field equations to a form in which a positive cosmological constant appears as a uniform density of gravitationally repulsive matter. Field equations with both positive and negative active gravitational mass densities of both primordial and continuously created matter incorporated along with two scalar fields coupled to geometry yield cosmological solutions that exhibit inflation, deceleration, coasting, acceleration, and a ‘big bounce’ instead of a ‘big bang’. The repulsive matter is identified as the back sides of the ‘drainholes’ introduced by the author in 1973 as solutions of those same field equations. Drainholes are topological tunnels in space which gravitationally attract on their front, entrance sides and repel more strongly on their back, exit sides. The front sides serve in part as the unseen particles of ‘dark matter’ needed to hold together the large scale structures seen in the universe, the back sides, as the misnamed ‘dark energy’ driving the accelerating expansion of the universe. Formation of cosmic voids, walls, filaments, and nodes is attributed to expulsion of drainhole entrances from voids made and populated by drainhole exits, and subsequent aggregation of the entrances on the void boundaries. All of these cosmological entities are found wrapped in one coherent package, namely, the field equations and the variational principle from which they are derived. The novelty of the underlying concepts gives rise to a number of issues and observations.

PACS numbers: PACS numbers: 98.80.Jk, 98.80.Cq, 95.35.+d, 95.36.+x

I. EINSTEIN’S UNJUSTIFIED ASSUMPTION

Albert Einstein, in his 1916 paper *Die Grundlage der allgemeinen Relativitätstheorie* [1] that gave a thorough presentation of the theory of gravity he had worked out over the preceding decade, made an assumption that does not hold up well under scrutiny. Stripped down to its barest form the assumption is that inertial mass, and by extension energy *via* $E = mc^2$, is a source of gravity and must therefore be coupled to the gravitational potential in the field equations of the general theory of relativity. The train of thought that brought him to this conclusion is described in §16, where he sought to extend his field equations for the vacuum, $\mathbf{R}_{\alpha\beta} - \frac{1}{2}\mathbf{R}g_{\alpha\beta} = 0$ as currently formulated, to include the contribution of a continuous distribution of gravitating matter of density μ , in analogy to the extension of the Laplace equation $\nabla^2\phi = 0$ for the newtonian gravitational potential ϕ to the Poisson equation $\nabla^2\phi = 4\pi\kappa\mu$, where κ is Newton’s gravitational constant. Einstein referred to μ as the “density of matter”, without specifying what was meant by ‘matter’ or its ‘density’. Invoking the special theory’s identification of “inert mass” with “energy, which finds its complete mathematical expression in . . . the energy-tensor”, he concluded that “we must introduce a corresponding energy-tensor of matter $\mathbf{T}_\sigma^\alpha$ ”. Further describing this energy-tensor as “corresponding to the density μ in Poisson’s equation”, he arrived at the extended field equa-

tions $\mathbf{R}_{\alpha\beta} - \frac{1}{2}\mathbf{R}g_{\alpha\beta} = \frac{8\pi\kappa}{c^2}T_{\alpha\beta}$, in which, for a “frictionless adiabatic fluid” of “density” μ , pressure p (a form of kinetic energy), and proper four-velocity distribution u^α , he took $T^{\alpha\beta}$ to be $\mu u^\alpha u^\beta + (p/c^2)(u^\alpha u^\beta - g^{\alpha\beta})$.

Clearly, Einstein’s procedure fails to distinguish between the ‘active gravitational mass’ of matter, which measures how much gravity it produces and is the sole contributor to the “density of matter” in Poisson’s equation, and the “inert mass” of matter, which measures how much it accelerates in response to forces applied to it, an effect entirely different from the production of gravity. These two conceptually different masses, along with yet a third, all occur in Newton’s gravitational equation

$$m_i a_B = F_{AB} = -\kappa \frac{m_p M_a}{r^2}, \quad (1)$$

in which M_a is the *active* gravitational mass of a gravitating body A, m_i is the *inertial* (“inert”) mass of a body B being acted upon by the gravity of A, and m_p is the *passive* gravitational mass of B, a measure of the strength of B’s ‘sensing’ of the gravitational field around A. That in suitable units $m_i = m_p$ for all bodies is another way of saying that all bodies respond with the same accelerations to the same gravitational fields, that, in consequence, the notion of a ‘gravitational force’ is irrelevant, but the notion of a ‘gravitational field’ is not. Simple thought experiments of Galileo (large stone and smaller stone tied together) [2] and Einstein (body suspended by rope in elevator) [3] make it clear that bodies do all respond alike — an observation now treated as a principle, the (weak) ‘principle of equivalence’, experimentally, if somewhat redundantly, well confirmed. That this passive-inertial mass has any relation to active gravita-

*URL: <http://euclid.colorado.edu/~ellis>; Electronic address: ellis@euclid.colorado.edu

tional mass is not apparent in Eq. (1), where, unlike m_i and m_p , M_a represents a property of A, not of B. But Newton's equation for the gravitational action of B on A reads

$$M_i a_A = F_{BA} = -\kappa \frac{M_p m_a}{r^2}. \quad (2)$$

Application of Newton's law of action and reaction allows the inference that F_{AB} and F_{BA} have the same magnitude, from which follows that $m_a/m_p = M_a/M_p$, hence that the ratio of active gravitational mass to passive gravitational mass, thus to inertial mass, is the same for all bodies. It would seem likely that Einstein relied, either consciously or unconsciously, on this consequence of Newton's laws when he assumed that "inert mass" should contribute to the "density of matter" as a source of gravity in the field equations.

Newton's law of action and reaction is applicable to the bodies A and B only under the condition that gravity acts at a distance instantaneously, that is, at infinite propagation speed. But the general theory of relativity that Einstein was expounding is a field theory in which gravitational effects propagate at finite speed. Within his own theory of gravity there is, therefore, no obvious justification for Einstein's assumption that inertial mass (and therefore energy) is equivalent to active gravitational mass. This, however, is not to say that there is no relation at all between the two kinds of mass. There is, for example, the seemingly universal coincidence that wherever there is matter made of atoms there are to be found both inertial mass and active gravitational mass. Indeed, the fact that Newton's theory is a close first approximation to Einstein's would argue for some proportionality between m_a and m_p for such matter in bulk — not, however, for each individual constituent of such matter. A 1986 analysis of lunar data concluded that the ratio of m_a to m_p for aluminum differs from that for iron by less than 4×10^{-12} [4]. An earlier, Cavendish balance experiment had put a limit of 5×10^{-5} on the difference of these ratios for bromine and fluorine [5]. But those results are only for matter in bulk, that is, matter made of atoms and molecules. It is entirely possible that electrons, for example, do not gravitate at all, for no one has ever established by direct observation that they do, nor is it likely that anyone will. There is in the literature an argument that purports to show that if the ratio m_a/m_p is the same for two species of bulk matter, then electrons must be generators of gravity [6], but that argument can be seen on careful examination to rest on an unrecognized, hidden assumption, namely that, in simplest form, the gravitational field of a hydrogen atom at a distance could be distinguished from that of a neutron at the same distance — another assumption no one has tested or is likely to test by direct observation.

Einstein's assumption that energy and inertial mass are sources of gravity has survived to the present virtually unchallenged. It has generated a number of consequences that have directed much of the subsequent re-

search in gravitation theory — indeed, misdirected it if his assumption is wrong. Among them are the following:

- The impossibility, according to Penrose–Hawking singularity theorems, of avoiding singularities in the geometry of space-time without invoking 'negative energy', which is really just energy coupled to gravity with polarity opposite to that of the coupling of matter to gravity.
- The presumption that the extra, fifth dimension in Kaluza–Klein theory must be a spatial dimension rather than a dimension of another type.
- The belief that all the extra dimensions in higher-dimensional theories must be spatial, causing the expenditure of much effort in mental gymnastics to explain why they are not apparent to our senses in the way that the familiar three spatial dimensions are.

Denying Einstein's assumption relieves one of the burden of these troublesome conclusions and opens the door to other, more realistic ones.

II. NEW, IMPROVED FIELD EQUATIONS

If Einstein's assumption is to be disallowed, then his source tensor for a continuous distribution of gravitating matter, $T^{\alpha\beta} = \mu u^\alpha u^\beta + (p/c^2)(u^\alpha u^\beta - g^{\alpha\beta})$, must be modified or replaced. One might think to simply drop the second term and take $T^{\alpha\beta} = \mu u^\alpha u^\beta$, the energy-momentum tensor of the matter. This would be inconsistent, for the μ in that tensor is the density of inertial-passive mass, which we are now not assuming to be the same as active gravitational mass. What to do instead?

At the same time that Einstein was creating his field equations, Hilbert was deriving the field equations for (in particular) empty space from the variational principle $\delta \int \mathbf{R} |g|^{\frac{1}{2}} d^4x = 0$ [7]. This is the most straightforward extension to the general relativity setting of the variational principle $\delta \int |\nabla\phi|^2 d^3x = 0$, whose Euler–Lagrange equation is equivalent to the Laplace equation $\nabla^2\phi = 0$ for the newtonian potential ϕ . Modifying that principle to $\delta \int (|\nabla\phi|^2 + 8\pi\kappa\mu\phi) d^3x = 0$ generates the Poisson equation $\nabla^2\phi = 4\pi\kappa\mu$. The most straightforward extension of this modified principle to general relativity is

$$\delta \int (\mathbf{R} - \frac{8\pi\kappa}{c^2}\mu) |g|^{\frac{1}{2}} d^4x = 0, \quad (3)$$

for which the Euler–Lagrange equations are equivalent to

$$\mathbf{R}_{\alpha\beta} - \frac{1}{2}\mathbf{R}g_{\alpha\beta} = -\frac{4\pi\kappa}{c^2}\mu g_{\alpha\beta}, \quad (4)$$

which makes $T_{\alpha\beta} = -\frac{1}{2}\mu g_{\alpha\beta}$, with μ now the *active* gravitational mass density, as it should be. Equivalent to this equation is $\mathbf{R}_{\alpha\beta} = \frac{4\pi\kappa}{c^2}\mu g_{\alpha\beta}$, the 00 component

of which reduces in the slowly varying, weak field approximation precisely to the Poisson equation. The vanishing of the divergence of the Einstein tensor field on the lefthand side of Eq. (4) entails that $0 = T_{\alpha}^{\beta}{}_{;\beta} = -\frac{1}{2}(\mu_{;\beta}g_{\alpha}^{\beta} + \mu g_{\alpha}^{\beta}{}_{;\beta}) = -\frac{1}{2}\mu_{;\alpha}$, hence that μ is a constant. This would seem to be a comedown from the equations of motion of the matter distribution implied by the vanishing of the divergence of Einstein's $T^{\alpha\beta}$, but those equations are unrealistic in that they have the density of active mass playing the role that properly belongs to the density of inertial mass as the coefficient of the four-acceleration of the matter. As will be seen, the new, improved field equations to be arrived at will allow μ to vary.

To widen the range of space-time geometries admitted by the field equations one can in the usual way add to the action integrand of Eq. (3) terms related to such things as scalar fields and electromagnetic fields. In particular, one can add a cosmological constant term, changing the integrand to $\mathbf{R} - \frac{8\pi\kappa}{c^2}\mu + 2\Lambda$ and the field equations to

$$\mathbf{R}_{\alpha\beta} - \frac{1}{2}\mathbf{R}g_{\alpha\beta} = -\frac{4\pi\kappa}{c^2}(\mu + \bar{\mu})g_{\alpha\beta}, \quad (5)$$

where $\frac{4\pi\kappa}{c^2}\bar{\mu} = -\Lambda$. A positive cosmological constant Λ thus appears in this context to be a (mis)representation of a negative active mass density $\bar{\mu}$ of a continuous distribution of gravitationally repulsive matter. An excess of this negative density over the positive active mass density μ of attractive matter could drive an accelerating cosmic expansion, and in doing so would provide a solution to the vexing ‘Cosmological Constant Problem’. Leaving aside for the moment the question of how such a negative mass density might arise, let us explore the consequences of presuming one exists, by studying cosmological solutions of field equations that incorporate a positive mass density μ , a negative mass density $\bar{\mu}$, and minimally coupled scalar fields ϕ (not the newtonian ϕ) and ψ . The variational principle

$$\delta \int [\mathbf{R} - \frac{8\pi\kappa}{c^2}(\mu + \bar{\mu}) + 2\phi^{\cdot\gamma}\phi_{\cdot\gamma} - 2\psi^{\cdot\gamma}\psi_{\cdot\gamma}] |g|^{\frac{1}{2}} d^4x = 0 \quad (6)$$

combines these elements and generates the field equations

$$\mathbf{R}_{\alpha\beta} - \frac{1}{2}\mathbf{R}g_{\alpha\beta} = \frac{8\pi\kappa}{c^2}T_{\alpha\beta} := -\frac{4\pi\kappa}{c^2}(\mu + \bar{\mu})g_{\alpha\beta} - 2(\phi_{\cdot\alpha}\phi_{\cdot\beta} - \frac{1}{2}\phi^{\cdot\gamma}\phi_{\cdot\gamma}g_{\alpha\beta}) + 2(\psi_{\cdot\alpha}\psi_{\cdot\beta} - \frac{1}{2}\psi^{\cdot\gamma}\psi_{\cdot\gamma}g_{\alpha\beta}), \quad (7)$$

and

$$2(\Box\phi)\phi_{\cdot\alpha} - 2(\Box\psi)\psi_{\cdot\alpha} := 2\phi^{\cdot\gamma}{}_{;\gamma}\phi_{\cdot\alpha} - 2\psi^{\cdot\gamma}{}_{;\gamma}\psi_{\cdot\alpha} = -\frac{4\pi\kappa}{c^2}(\mu + \bar{\mu})_{\cdot\alpha}, \quad (8)$$

the latter of which is a derivation from the former, corresponding to the identity $T_{\alpha}^{\beta}{}_{;\beta} = 0$. In this variational principle only the metric components $g_{\alpha\beta}$ are varied. A superficial reason for disallowing variation of ϕ and ψ is that varying them would produce the conclusion that $\Box\phi = 0$ and $\Box\psi = 0$, thus, in light of Eq. (8), the unwanted restriction that $\mu + \bar{\mu}$ is constant. A more fundamental reason for not varying them will be discussed later. Notice that the presence of two scalar fields allows one to disregard by circumvention the question of whether the coupling of a scalar field to the space-time geometry should have positive polarity or negative polarity. Indeed, both polarities are consistent with Einstein's assumption as well as with its denial, inasmuch as the ‘energy’ of a scalar field χ included in $\chi_{\cdot\alpha}\chi_{\cdot\beta} - \frac{1}{2}\chi^{\cdot\gamma}\chi_{\cdot\gamma}g_{\alpha\beta}$ is of a nature entirely different from that of the kinetic energy in Einstein's “energy-tensor”, thus should have no connection to that tensor.

III. COSMIC EVOLUTION EQUATIONS

For a Robertson–Walker metric $c^2 dt^2 - R^2(t)ds^2$ (with t in seconds, s dimensionless, c in centimeters/second,

and R in centimeters) and dimensionless scalar fields $\phi = \alpha(t)$ and $\psi = \beta(t)$ the field equations (7) reduce to

$$3\frac{\dot{R}^2/c^2 + k}{R^2} = -\frac{4\pi\kappa}{c^2}(\mu + \bar{\mu}) - \frac{\dot{\alpha}^2 - \dot{\beta}^2}{c^2}, \quad (9)$$

and

$$\frac{2}{c^2}\frac{\ddot{R}}{R} + \frac{\dot{R}^2/c^2 + k}{R^2} = -\frac{4\pi\kappa}{c^2}(\mu + \bar{\mu}) + \frac{\dot{\alpha}^2 - \dot{\beta}^2}{c^2}, \quad (10)$$

where $k = -1, 0$, or 1 , the uniform curvature of the spatial metric ds^2 . These equations, which are replacements for the well-studied Friedmann–Lemaître cosmological equations, are equivalent together to

$$\frac{1}{c^2}\frac{\ddot{R}}{R} = -\frac{4\pi\kappa}{3c^2}(\mu + \bar{\mu}) - \frac{k}{R^2} - \frac{\dot{\alpha}^2 - \dot{\beta}^2}{3c^2} \quad (11)$$

and

$$\frac{1}{c^2}\frac{\ddot{R}}{R} = -\frac{4\pi\kappa}{3c^2}(\mu + \bar{\mu}) + \frac{2(\dot{\alpha}^2 - \dot{\beta}^2)}{3c^2}. \quad (12)$$

Equations (8) reduce to

$$\frac{2}{c^2} \left(\ddot{\alpha} + 3 \frac{\dot{R}}{R} \dot{\alpha} \right) \dot{\alpha} - \frac{2}{c^2} \left(\ddot{\beta} + 3 \frac{\dot{R}}{R} \dot{\beta} \right) \dot{\beta} = -\frac{4\pi\kappa}{c^2} \partial_t(\mu + \bar{\mu}) \quad (13)$$

for the time component and $\partial_\alpha(\mu + \bar{\mu}) = 0$ for the space components. Equation (13) is equivalent to

$$\frac{1}{c^2} \partial_t \left(R^6 (\dot{\alpha}^2 - \dot{\beta}^2) \right) = -\frac{4\pi\kappa}{c^2} R^6 \partial_t(\mu + \bar{\mu}). \quad (14)$$

To give substance to the densities μ and $\bar{\mu}$, let us invoke two kinds of gravitating matter: a) *primordial matter*, existent at all times, never changing in amount, with its net density $\mu + \bar{\mu}$ inversely proportional to the cube of the scale factor R ; and b) *continuously created* (or *destroyed*) *matter*, coming steadily into existence (or passing out of existence) at a rate just sufficient to keep its net density constant. For primordial matter we have

$$\mu_P(t) + \bar{\mu}_P(t) = (\mu_{P,0} + \bar{\mu}_{P,0}) \frac{R^3(t_0)}{R^3(t)}, \quad (15)$$

where t_0 denotes the value of t at the present epoch, and $\mu_{P,0}$ and $\bar{\mu}_{P,0}$ are the present values of the primordial densities μ_P and $\bar{\mu}_P$. For continuously created matter we have

$$\mu_C(t) + \bar{\mu}_C(t) = \mu_{C,0} + \bar{\mu}_{C,0}, \quad (16)$$

where $\mu_{C,0}$ and $\bar{\mu}_{C,0}$ are the present values of the continuously created densities μ_C and $\bar{\mu}_C$. Under the assumption that the net densities $\mu_P + \bar{\mu}_P$ and $\mu_C + \bar{\mu}_C$ are additive, Eq. (14) turns into

$$\frac{1}{c^2} \partial_t \left(R^6 (\dot{\alpha}^2 - \dot{\beta}^2) \right) = \frac{4\pi\kappa}{c^2} (\mu_{P,0} + \bar{\mu}_{P,0}) R^3(t_0) \partial_t(R^3), \quad (17)$$

which integrates to

$$\frac{1}{c^2} (\dot{\alpha}^2 - \dot{\beta}^2) = \frac{4\pi\kappa}{c^2} (\mu_{P,0} + \bar{\mu}_{P,0}) \frac{R^3(t_0)}{R^3} + \frac{B}{R^6}, \quad (18)$$

where B is the constant of integration, with units cm^4 .

At this point it is convenient to set

$$A_P := -\frac{4\pi\kappa}{c^2} (\mu_{P,0} + \bar{\mu}_{P,0}) R^3(t_0) \quad (19)$$

(units cm) and

$$A_C := -\frac{4\pi\kappa}{c^2} (\mu_{C,0} + \bar{\mu}_{C,0}) \quad (20)$$

(units cm^{-2}). Equation (18) becomes

$$\frac{1}{c^2} (\dot{\alpha}^2 - \dot{\beta}^2) = \frac{B - A_P R^3}{R^6}, \quad (21)$$

and substitution from Eqs. (19), (20), and (21) into Eqs. (11) and (12) produces

$$\frac{1}{c^2} \frac{\dot{R}^2}{R^2} = \frac{P_1(R)}{3R^6} = \frac{A_C}{3} - \frac{k}{R^2} + \frac{2A_P}{3R^3} - \frac{B}{3R^6} \quad (22)$$

and

$$\frac{1}{c^2} \frac{\ddot{R}}{R} = \frac{P_2(R)}{3R^6} = \frac{A_C}{3} - \frac{A_P}{3R^3} + \frac{2B}{3R^6}, \quad (23)$$

where

$$P_1(R) := A_C R^6 - 3kR^4 + 2A_P R^3 - B \quad (24)$$

and

$$P_2(R) := A_C R^6 - A_P R^3 + 2B. \quad (25)$$

According to Eqs. (19) and (20), positive values for A_P and A_C equate to negative net densities for primordial matter and continuously created matter, signifying that on balance each kind gravitationally repels all other matter more strongly than it attracts other matter. This excess of repulsion over attraction will be presumed to exist for both types. With thus $A_P > 0$ and $A_C > 0$ assumed, along with an added assumption that $B > 0$, several properties of the scale factor R as a solution of these equations can be inferred rather easily, to wit:

- For each of $k = -1, 0, 1$ there is a positive number $R_{\min}$ (the least positive root of the polynomial $P_1(R)$, where $\dot{R} = 0$) that Eq. (22) does not allow R to go below. (See Fig. 1.) This sets a positive lower bound on the compression of space and thereby rules out the development of a ‘big bang’ singularity.
- For $k = 1$ and some choices of A_C , A_P , and B there are in addition to $R_{\min}$ other values $R_{\max}$ and $R_{\min}^*$ of R at which P_1 , and consequently $\dot{R}$, vanish.
- Wherever $P_1(R) \leq 0$, $P_2(R) \geq R^3(A_C R^3 - 2kR + A_P) = \frac{1}{2} R P_1'(R)$. Because, with one exception (the graph labeled b in Fig. 1), $P_1'(R_{\min}) > 0$, also $P_2(R_{\min}) > 0$, and therefore $\ddot{R}$ is positive when $R = R_{\min}$. This tells that in place of a ‘big bang’ there is a ‘bounce’ off a state of maximum compression at the time when $R = R_{\min}$. The lone exception has $k = 1$, $P_1'(R_{\min}) = 0$, and $R(t) = R_{\min} = R_{\max}$ for all time, modeling a static, spherical universe of radius $R_{\min}$.
- With the time $t = 0$ chosen so that $R(0) = R_{\min}$, $R(t)$ is symmetric about $t = 0$, the evolution of the universe after time zero thus being mirrored in reverse as a devolution of the universe before time zero.




FIG. 1: Graphs of $P_1(R)$ for $k = -1, 0$, and 1 , and generic positive values of the parameters A_C, A_P , and B . Values of R for which $P_1(R) < 0$ are excluded from the range of $R(t)$ by Eq. (22).

- Whereas for $k = -1$ and $k = 0$ (Fig. 1), and $k = 1$ (Fig. 1, graph a), the universe expands from $R_{\min}$ to infinity as $t \rightarrow \pm\infty$, other behaviors are possible

when $k = 1$, most notably: (Fig. 1, graph b) static spherical universe as noted above; (Fig. 1, graph c) spherical universe oscillating for all time between minimum radius $R_{\min}$ and maximum radius $R_{\max}$.

- The polynomial $P_2(R)$ is quadratic in R^3 , with discriminant $A_P^2 - 8A_CB$. If $B > A_P^2/8A_C$, then $P_2(R)$ has no positive root, so $\ddot{R} > 0$ at all times, thus the universal expansion described by $R(t)$ for times after $t = 0$ is always accelerating. If $B = A_P^2/8A_C$, then $P_2(R)$ has a single positive root, so the expansion is, except for a momentary pause, accelerating at all positive times, unless, as in Fig. 1, graph b, $R(t) = R_{\min} = R_{\max}$ or, as in Fig. 1, graph d, $R(t) = R_{\min}^* = R_{\max}$. If $B < A_P^2/8A_C$, then $P_2(R)$ has two positive roots,

$$R_d := \left(\frac{A_P}{2A_C} - \frac{\sqrt{A_P^2 - 8A_CB}}{2A_C} \right)^{\frac{1}{3}}, \quad (26)$$

which marks a transition from the initial accelerating expansion associated with the bounce at $t = 0$ to an interval of decelerating expansion, and

$$R_a := \left(\frac{A_P}{2A_C} + \frac{\sqrt{A_P^2 - 8A_CB}}{2A_C} \right)^{\frac{1}{3}}, \quad (27)$$

which marks a return to accelerating expansion (if the expansion is destined to continue forever, as opposed to some of the other possible behaviors when $k = 1$).

- For the infinitely expanding models the ‘Hubble parameter’ H and the ‘acceleration parameter’ Q behave asymptotically as follows:

$$H := \dot{R}/R, \quad \frac{1}{c^2} H^2 = \frac{A_C}{3} - \frac{3kR^4 - 2A_P R^3 + B}{3R^6} \rightarrow \frac{A_C}{3} \quad \left\{ \begin{array}{l} \text{from below if } k > 0 \\ \text{from above if } k \leq 0 \end{array} \right\} \text{ as } R \rightarrow \infty, \quad (28)$$

and

$$Q := \frac{\ddot{R}/R}{(\dot{R}/R)^2}, \quad Q = c^2 \frac{A_C R^6 - A_P R^3 + 2B}{3H^2 R^6} \quad (29)$$

$$= 1 + c^2 \frac{kR^4 - A_P R^3 + B}{H^2 R^6} \rightarrow 1 \quad \left\{ \begin{array}{l} \text{from above if } k > 0 \\ \text{from below if } k \leq 0 \end{array} \right\} \text{ as } R \rightarrow \infty. \quad (30)$$

Either of these entails that $R(t) \sim C e^{\pm \sqrt{A_C/3} c t}$, for some constant C , as $t \rightarrow \pm\infty$.

- For $k = -1$ or 0 , and for some cases of $k = 1$, H has a maximum value $H_{\max}$ at $R = R_{H_{\max}}$, where




FIG. 2: Graphs of $H(R)$ and $Q(R)$ for generic positive values of the parameters A_C , A_P , and B , showing early stage inflation followed by a decline in H to its asymptotic limit $H(\infty) = c\sqrt{A_C/3}$, and (for $B < A_P^2/8A_C$) transitions of Q from inflationary acceleration to deceleration at R_d and back to acceleration at R_a to the asymptotic limit $Q(\infty) = 1$.

$dH/dR = c^2(kR^4 - A_P R^3 + B)/HR^7 = 0$. If $k = 0$, $R_{H_{\max}} = \sqrt[3]{B/A_P}$ and $H_{\max} = c\sqrt{A_C/3 + A_P^2/3B}$. As seen in Fig. 2, H rises sharply from 0 at $R_{\min}$ to $H_{\max}$ at $R_{H_{\max}}$, then reverses and tails off asymptotically to $c\sqrt{A_C/3}$. One can show that $R_{\min} \sim \sqrt[3]{B/2A_P}$, $R_{H_{\max}} \sim \sqrt[3]{B/A_P}$, $R_d \sim \sqrt[3]{2B/A_P}$, and $H_{\max} \sim cA_P/\sqrt{3B}$ as $B \rightarrow 0$ with A_C and A_P fixed. Thus, as $B \rightarrow 0$ with A_C and A_P fixed, $R_{\min}$, $R_{H_{\max}}$, and R_d are squeezed together closer and closer to 0, and $H_{\max}$ grows without bound. This clearly is a recipe for an explosive postbounce inflation ending with a ‘graceful’ exit initiated by the onset of deceleration when R reaches R_d .

- As Fig. 2 exhibits, for $k = -1$, $k = 0$, and $k = 1$, graph a, the acceleration parameter Q , descending from ∞ at $R_{\min}$, passes through 1 at $R_{H_{\max}}$, bottoms out with a minimum value $Q_{\min}$ at $R_{Q_{\min}}$, where $dQ/dR = 0$, then creeps slowly back to 1 as $R \rightarrow \infty$ (with a late pass through 1 if $k = 1$). When $k = 0$, $Q_{\min} = -\frac{1}{2} + \frac{3}{2}\sqrt{A_C B/(A_P^2 + A_C B)}$ at




FIG. 3: Graphs of $H(R)$ and $R(t)$ for $k = 1$ and generic positive values of the parameters A_C , A_P , and B associated with graph c of Fig. 1, showing repetitive, identical periods of expansion and contraction, each beginning with a stage of rapid inflation from a bounce at $R = R_{\min}$, which is followed by a less rapid expansion to $R = R_{\max}$, then a mirror-image contraction to an ending stage of rapid deflation into the next bounce at $R = R_{\min}$. The functions are related by $\dot{R}(t)/R(t) =: H(R(t))$.

$R_{Q_{\min}} = \left(B/A_P + \sqrt{B^2/A_P^2 + B/A_C}\right)^{\frac{1}{3}}$, which, with A_C and A_P fixed, is asymptotic to $\sqrt[6]{B/A_C}$ as $B \rightarrow 0$.

- If $k = 1$ and A_C , A_P , and B are such as to generate graph c of Fig. 1, then for the section bounded by $R_{\min}$ and $R_{\max}$ the behavior of R is cyclical, as Fig. 3 portrays graphically.

IV. COSMOLOGICAL SOLUTIONS

A. Flat open universe ($k = 0$)

When $k = 0$, so that space is perfectly flat, $R_{\min}^3 = -A_P/A_C + \sqrt{A_P^2 + A_C B/A_C}$ and it is straightforward to integrate Eqs. (22) and (23) (which is equivalent to a




FIG. 4: Graph of $R(t)$ for $k = 0$ and generic positive values of A_C , A_P , and B with $B < A_P^2/8A_C$, showing early acceleration from $t = 0$ to $t = t_d$, deceleration from t_d to t_a , and ultimately exponential acceleration after t_a ; truncated graph of $R(t)$ with approximators $\hat{R}(t)$ and $\tilde{R}(t)$ as identified in Eqs. (34) and (36).

readily integrable equation linear in R^3), with the result that

$$R^3(t) = \left(R_{\min}^3 + \frac{A_P}{A_C} \right) \cosh \left(\sqrt{3A_C} ct \right) - \frac{A_P}{A_C}. \quad (31)$$

One can show that if $B \leq A_P^2/8A_C$, so that R_d and R_a exist, then $R_{\min} < R_d \leq R_a$. The postbounce times t_d and t_a at which $R = R_d$ and $R = R_a$ are the positive solutions of

$$\cosh \left(\sqrt{3A_C} ct \right) = \frac{R_{\{d,a\}}^3 + A_P/A_C}{R_{\min}^3 + A_P/A_C}. \quad (32)$$

Figure 4 displays the evolution of $R(t)$ for generic values of A_C , A_P , and B with $B < A_P^2/8A_C$, showing the acceleration during the interval from $t = 0$ to $t = t_d$, the deceleration during the interval from t_d to t_a , and the ultimately exponential acceleration after t_a .

For times $t \approx 0$, Eq. (31) yields the approximation

$$R^3(t) \approx R_{\min}^3 \left(1 + \frac{3A_C}{2} \frac{R_{\min}^3 + A_P/A_C}{R_{\min}^3} c^2 t^2 \right), \quad (33)$$

which produces the further approximation that

$$R(t) \approx \hat{R}(t) := R_{\min} + \frac{A_C}{2} \left(R_{\min} + \frac{A_P}{A_C} \frac{1}{R_{\min}^2} \right) c^2 t^2, \quad (34)$$

thus that $R(t)$ is approximately parabolic, opening upward from a vertex at $R_{\min}$, with an inflationary ‘steepness’ factor that grows as $c^2 A_P / 2R_{\min}^2$ as $R_{\min} \rightarrow 0$.

Equivalent to Eq. (33) is

$$R^3(t) - R_{\min}^3 \approx \frac{3A_C}{2} \left(R_{\min}^3 + \frac{A_P}{A_C} \right) (ct)^2. \quad (35)$$

After $R(t)$ has inflated to the point that $R_{\min}/R(t) \ll 1$, $(R^3(t) - R_{\min}^3)^{1/3}$ will differ but little from $R(t)$, and then, for so long as t remains small enough for the approximate Eq. (33) to hold, the approximation

$$R(t) \approx \tilde{R}(t) := \left[\frac{3A_C}{2} \left(R_{\min}^3 + \frac{A_P}{A_C} \right) \right]^{1/3} (ct)^{2/3} \quad (36)$$

will be valid. Figure 4 shows the approximations to $R(t)$.

B. Nonflat open ($k = -1$) and closed ($k = 1$) universes

When $k = -1$ or 1 , elementary symbolic solutions of Eqs. (22) and (23) are unavailable, so one must resort to numerical solution of Eq. (23). The integration proceeds most smoothly when carried out for the equivalent equation

$$\ddot{U} = 3c^2 A_C U - \frac{6c^2 k}{R_{\min}^2} U^{1/3} + \frac{3c^2 A_P}{R_{\min}^3}, \quad (37)$$

where $U := S^3$ and $S := R/R_{\min}$, with initial conditions $U(0) = 1$ and $\dot{U}(0) = 0$, obtained from $S(0) = R(0)/R_{\min} = R_{\min}/R_{\min} = 1$ and $\dot{S}(0) = \dot{R}(0)/R_{\min} = 0$. (Note that this equation and the initial conditions apply equally well when $k = 0$.)

To enable the integration, numerical values must be chosen for the parameters other than k that appear in Eq. (37), namely, A_C , A_P , and B (implicitly through $R_{\min}$). For this determination let us impose the requirement that the solution(s) provide good fits to a Hubble diagram that incorporates the data from recent supernovae observations. Toward that end a number of definitions are in order:

- t_0 , the time of the present epoch, i. e., the time elapsed since the bounce at $t = 0$, in seconds.
- $H_0 := H(t_0)$, the present value of the Hubble parameter.

- $R_0 := c/H_0$, usually referred to as the ‘Hubble radius’ and as the ‘radius of the visible universe’.
- $z := R(t_0)/R - 1$, the redshift function.
- $z_d := R(t_0)/R_d - 1$ and $z_a := R(t_0)/R_a - 1$, the redshifts associated with the transitions from acceleration to deceleration and from deceleration back to acceleration.
- $\tilde{R} := R/R(t_0) = 1/(1+z)$, a conventional notation.
- $\tilde{R}_{\min} := R_{\min}/R(t_0)$, and $\tilde{R}_{\{d,a\}} := R_{\{d,a\}}/R(t_0)$.
- $\lambda := c/H_0 R(t_0) = R_0/R(t_0)$, a useful parameter.
-

$$\Omega_C := \frac{c^2}{3H_0^2} A_C, \quad (38)$$

$$\Omega_k := -\frac{c^2}{H_0^2 R^2(t_0)} k = -\lambda^2 k, \quad (39)$$

$$\Omega_P := \frac{2c^2}{3H_0^2 R^3(t_0)} A_P = \frac{2H_0 \lambda^3}{3c} A_P, \quad (40)$$

and

$$\Omega_B := -\frac{c^2}{3H_0^2 R^6(t_0)} B = -\frac{H_0^4 \lambda^6}{3c^4} B. \quad (41)$$

With these definitions Eq. (22) is equivalent to

$$H^2 = H_0^2 \left(\Omega_C + \Omega_k \frac{1}{\tilde{R}^2} + \Omega_P \frac{1}{\tilde{R}^3} + \Omega_B \frac{1}{\tilde{R}^6} \right), \quad (42)$$

which upon evaluation at t_0 yields

$$\Omega_C + \Omega_k + \Omega_P + \Omega_B = 1. \quad (43)$$

On the other hand, Eq. (23) is equivalent to

$$\frac{\ddot{R}}{\tilde{R}} = H_0^2 \left(\Omega_C - \frac{1}{2} \Omega_P \frac{1}{\tilde{R}^3} - 2 \Omega_B \frac{1}{\tilde{R}^6} \right) \quad (44)$$

$$= H_0^2 \frac{2 \Omega_C \tilde{R}^6 - \Omega_P \tilde{R}^3 - 4 \Omega_B}{2 \tilde{R}^6}, \quad (45)$$

which yields, as roots of $\ddot{R} = 0$,

$$\tilde{R}_d := \left(\frac{\Omega_P}{2 \Omega_C} - \frac{\sqrt{\Omega_P^2 + 8 \Omega_C \Omega_B}}{2 \Omega_C} \right)^{\frac{1}{3}}, \quad (46)$$

and

$$\tilde{R}_a := \left(\frac{\Omega_P}{2 \Omega_C} + \frac{\sqrt{\Omega_P^2 + 8 \Omega_C \Omega_B}}{2 \Omega_C} \right)^{\frac{1}{3}} \quad (47)$$

(obtainable also from Eqs. (26) and (27) by substitution from Eqs. (38), (40), and (41)).

To produce a curve to match against the data points of a Hubble diagram one needs the following standard formula for the luminosity distance D_L of a photon-emitting astronomical object at redshift z :

$$D_L(z) = (1+z) R(t_0) r_k \left(\frac{c}{R(t_0)} \int_0^z \frac{1}{\dot{H}^*(u)} du \right). \quad (48)$$

Here

$$r_k(\rho) := \begin{cases} \sinh \rho & \text{if } k = -1, \\ \rho & \text{if } k = 0, \\ \sin \rho & \text{if } k = 1, \end{cases} \quad (49)$$

and

$$\dot{H}^*(z) := H_0 [\Omega_C + \Omega_k (1+z)^2 + \Omega_P (1+z)^3 + \Omega_B (1+z)^6]^{\frac{1}{2}}. \quad (50)$$

In terms of the parameter λ the luminosity distance formula reads

$$D_L(z) = (1+z) \frac{c}{\lambda H_0} r_k \left(\lambda \int_0^z \frac{du}{[\Omega_C + \Omega_k (1+u)^2 + \Omega_P (1+u)^3 + \Omega_B (1+u)^6]^{\frac{1}{2}}} \right). \quad (51)$$

From D_L one constructs the distance modulus μ , the difference between the apparent magnitude and the absolute magnitude of the object in question at redshift z ;

this reduces in the usual way to

$$\mu(z) = 5 \log_{10} \left(\frac{D_L(z)}{10 \text{ pc}} \right) = 25 + \log_{10} \left(\frac{D_L(z)}{1 \text{ Mpc}} \right). \quad (52)$$

From $\tilde{R} = 1/(1+z)$ follows $-\dot{z}/(1+z) = \dot{\tilde{R}}/\tilde{R} = \dot{H}^*(z)$, which upon integration yields the following formula for

the time $T(z)$ elapsed between an event occurring at redshift z and the present time t_0 (when $z = 0$):

$$T(z) = \int_0^z \frac{1}{(1+u)\dot{H}^*(u)} du = \frac{1}{H_0} \int_0^z \frac{du}{(1+u)[\Omega_C + \Omega_k(1+u)^2 + \Omega_P(1+u)^3 + \Omega_B(1+u)^6]^{\frac{1}{2}}}. \quad (53)$$

The distance modulus μ of Eq. (52) is the function to be fit to the relative magnitudes data obtained from observations of supernovae. The data to be used are those from the 182-member sample described in Sec. 3 of [8], provided at http://braeburn.pha.jhu.edu/~ariess/R06/sn_sample. The requisite calculations are carried out (using the indicated Mathematica functions) in two stages, the first of which comprises the following steps:

1. On first run, set $H_0 = 74.2 \text{ km s}^{-1}/\text{Mpc}$, the recent estimate of H_0 obtained by Riess, et al. [9]. On subsequent runs set $H_0 = 70.6 \text{ km s}^{-1}/\text{Mpc}$ and $H_0 = 77.8 \text{ km s}^{-1}/\text{Mpc}$, the extremes of the 4.8% confidence interval predicted in [9].
2. Set $B = 0$, thus $\Omega_B = 0$. This makes $R_d = R_{\min} = 0$, so will produce a ‘big bang’ model with $z_d = \infty$ and with no inflation (cf. Eq. (23), in which the last term, now absent, is the one that makes $\tilde{R}$ large when $R \approx 0$).
3. For $k = 0$, using NIntegrate as the integrator for $D_L(z)$ in Eq. (51), apply NonlinearModelFit to

$\mu(z)$ to obtain, subject to the constraint of Eq. (43), Chi-square best-fit values for Ω_P , Ω_C , and λ . Do the same for $k = -1$ and $k = 1$. (Note that when $k = 0$, λ disappears from Eq. (51), so is irrelevant to the fitting.)

4. From the best-fit values thus obtained, compute $\tilde{R}_a$ from Eq. (47), then z_a from $\tilde{R}_a = 1/(1+z_a)$.
5. From Eq. (53), by applying NDSolve to the initial-value problem $dT/dz = 1/(1+z)\dot{H}^*(z)$, $T(0) = 0$, compute $T(z)$ and then the implied time elapsed since the big bang as $\lim_{z \rightarrow \infty} T(z)$.
6. Compute the Chi-square statistic

$$\chi^2 := \sum_{i=1}^{182} \left(\frac{\mu(z_i) - \mu_i}{\sigma_i} \right)^2. \quad (54)$$

Table I presents the results of these computations.

TABLE I: Best-fit parameters (λ not shown) for ‘big bang’ model: B set to 0, thus $\Omega_B = 0$, so $R_d = R_{\min} = 0$ and $z_d = \infty$.

Inputs		Outputs						
H_0 (km s ⁻¹ /Mpc)	k	Ω_P	Ω_C	Ω_k	$\tilde{R}_a$	z_a	$\lim_{z \rightarrow \infty} T(z)$ (Gyr)	χ^2
70.6	-1	0.419	0.581	2.986×10^{-20}	0.712	0.405	12.14	163.075
	0	0.419	0.581	0	0.712	0.405	12.14	163.075
	1	0.419	0.581	-2.379×10^{-18}	0.712	0.405	12.14	163.075
74.2	-1	0.295	0.705	2.796×10^{-22}	0.594	0.684	12.77	160.829
	0	0.295	0.705	0	0.594	0.684	12.77	160.829
	1	0.295	0.705	-2.796×10^{-22}	0.594	0.684	12.77	160.829
77.8	-1	0.198	0.802	1.262×10^{-19}	0.498	1.008	13.57	183.225
	0	0.198	0.802	0	0.498	1.008	13.57	183.225
	1	0.198	0.802	-1.277×10^{-19}	0.498	1.008	13.57	183.225

To prepare for the second stage observe that Eqs. (46) and (47) imply that

$$\Omega_P = 2(\tilde{R}_d^3 + \tilde{R}_a^3)\Omega_C \quad \text{and} \quad \Omega_B = -\frac{1}{2}\tilde{R}_d^3\tilde{R}_a^3\Omega_C. \quad (55)$$

Equivalent to Eq. (43) is $\Omega_C + \Omega_P + \Omega_B = 1 - \Omega_k$, substitution into which from Eqs. (55), then back into Eqs. (55), yields

$$\Omega_C = \frac{1 - \Omega_k}{1 + 2(\tilde{R}_d^3 + \tilde{R}_a^3) - \frac{1}{2}\tilde{R}_d^3\tilde{R}_a^3}, \quad (56)$$

$$\Omega_P = \frac{2(\tilde{R}_d^3 + \tilde{R}_a^3)(1 - \Omega_k)}{1 + 2(\tilde{R}_d^3 + \tilde{R}_a^3) - \frac{1}{2}\tilde{R}_d^3\tilde{R}_a^3}, \quad (57)$$

and

$$\Omega_B = \frac{-\frac{1}{2}\tilde{R}_d^3\tilde{R}_a^3(1 - \Omega_k)}{1 + 2(\tilde{R}_d^3 + \tilde{R}_a^3) - \frac{1}{2}\tilde{R}_d^3\tilde{R}_a^3}, \quad (58)$$

where, from Eq. (39), $\Omega_k = -\lambda^2 k$.

Because the supernovae data extend into the past no further than $z = 1.755$, the best-fit parameters Ω_P , Ω_C , and λ will change by negligible amounts when Ω_B changes from 0 to a small nonzero number. For this reason a different strategy is called for in the second stage, as detailed in the following steps:

7. Set λ and z_a to the best-fit values found in the first stage (for $k = 0$, choose $\lambda = 1$). Assign z_d a large value.

8. From $\tilde{R} = 1/(1+z)$, compute $\tilde{R}_d$ and $\tilde{R}_a$ (reversing, in the latter case, the computation of z_a from $\tilde{R}_a$ in the first stage).

9. Using Eqs. (56), (57), (58), and $\Omega_k = -\lambda^2 k$, compute Ω_C , Ω_k , Ω_P , and Ω_B , and in turn $D_L(z)$ and $\mu(z)$ from Eqs. (51) and (52), applying NDSolve to the initial-value problem $dI/dz = H_0/\dot{H}(z)$, $I(0) = 0$ to compute the integral in Eq. (51).

10. Compute the Chi-square statistic and compare it to the corresponding number from the first stage. If it is appreciably larger, make z_d larger, thus R_d smaller and Ω_B closer to 0.

11. When a good match is found in Step 10, recover A_C , A_P , and B from Eqs. (38), (40), and (41), then $R_{\min}$ from $P_1(R_{\min}) = 0$ (cf. Eq. (24)) using NSolve.

With A_C , A_P , B and $R_{\min}$ in hand, proceed to the solution by NDSolve of Eq. (37) for $U(t)$ and $S(t) = [U(t)]^{\frac{1}{3}}$, with $U(0) = 1$ and $\dot{U}(0) = 0$, to obtain a full history of the model universe. As a check use FindRoot to get t_0 from $\dot{S}(t_0)/S(t_0) = H(t_0) = H_0$ and compare it to $T(z_{\min})$ from Eq. (53) (see Step 5), where $1 + z_{\min} = R(t_0)/R_{\min} = R_0/\lambda R_{\min}$. They must agree.

One can find t_{100} , the time at which R has doubled in size one hundred times from its minimum size at the bounce, by solving the equation $\log_2 S(t) = 100$ using FindRoot. If a different value for t_{100} is wanted, change z_d and repeat Steps 7–11 and the integration of Eq. (37).

Table II presents the results of these computations.

TABLE II: Best-fit parameters for ‘bounce’ model: $B > 0$, thus $\Omega_B < 0$ and $R_{\min} > 0$. (The columns for Ω_P , Ω_C , and Ω_k are identical to those of Table I, and are included by reference.)

Inputs					Outputs			
H_0 (km s ⁻¹ /Mpc)	k	λ	z_a	z_d	Ω_B	t_{100} (s)	t_0 (Gyr)	χ^2
70.6	-1	1.728×10^{-20}	0.405	1×10^{65}	-1.047×10^{-196}	1.02×10^{-35}	12.14	163.075
	0	1	0.405	1×10^{65}	-1.047×10^{-196}	1.02×10^{-35}	12.14	163.075
	1	1.543×10^{-9}	0.405	1×10^{65}	-1.047×10^{-196}	1.02×10^{-35}	12.14	163.075
74.2	-1	1.672×10^{-11}	0.684	1×10^{65}	-7.378×10^{-197}	1.15×10^{-35}	12.77	160.829
	0	1	0.684	1×10^{65}	-7.378×10^{-197}	1.15×10^{-35}	12.77	160.829
	1	1.672×10^{-11}	0.684	1×10^{65}	-7.378×10^{-197}	1.15×10^{-35}	12.77	160.829
77.8	-1	3.552×10^{-10}	1.008	1×10^{65}	-4.951×10^{-197}	1.34×10^{-35}	13.57	183.225
	0	1	1.008	1×10^{65}	-4.951×10^{-197}	1.34×10^{-35}	13.57	183.225
	1	3.573×10^{-10}	1.008	1×10^{65}	-4.951×10^{-197}	1.34×10^{-35}	13.57	183.225




FIG. 5: Best fit models of $\mu(z)$ for $k = 0$ (parameters shown in Tables II and I), plotted against the 182 SNe Ia data points they are fitted to. The corresponding plots for $k = -1$ and $k = 1$ are visually indistinguishable from these.

Figure 5 shows the best-fit models of $\mu(z)$ for $k = 0$, plotted against the 182 SNe Ia data points they are fitted to. The parameters are those of Table II. The corresponding plots for $k = -1$ and $k = 1$ are visually indistinguishable from the ones shown.

Figure 6 exhibits a complete postbounce history of the best-fit model for $H_0 = 74.2 \text{ km s}^{-1}/\text{Mpc}$ and $k = 0$. The graph is constructed from $R_{\min}^3 = -A_P/A_C + \sqrt{A_P^2 + A_C B}/A_C$, $S := R/R_{\min}$, and the equation, equivalent to Eq. (31),

$$R^3(t) = \frac{\sqrt{A_P^2 + A_C B}}{A_C} \cosh(\sqrt{3A_C} ct) - \frac{A_P}{A_C}, \quad (59)$$

with $A_P = 5.519 \times 10^{27} \text{ cm}$, $A_C = 1.360 \times 10^{-56} \text{ cm}^{-2}$, and $B = 5.348 \times 10^{-84} \text{ cm}^4$, obtained from the best-fit parameters Ω_P , Ω_C , and Ω_B by use of Eqs. (40), (38), and (41). The history comprises a brief interval of rapid inflation from the bounce, sliding into deceleration and a graceful exit into an era of coasting, followed by a return to ultimately exponential acceleration at $z = z_a = 0.684$ (when $t = 6.90 \text{ Gyr}$). Throughout the coasting era $Q(t) \approx Q_{\min} = -0.5 + 7.33 \times 10^{-98}$, reflective of the approximate proportionality $R(t) \propto (ct)^{2/3}$ implied by Eq. (36). At $t = t_0$, $Q(t) = 0.557$. Numerical integration of Eq. (37) produces the same graph for $k = 0$, and for

$k = \pm 1$ yields graphs visually indistinguishable from it.

V. DARK MATTER AND DARK ‘ENERGY’

A. Drainholes

For their success in modeling inflation, deceleration, and acceleration the cosmological solutions developed in Sec. IV depend on, among other modifications, the inclusion of the negative density $\bar{\mu}$ alongside the usual positive density μ in the variational principle of Eq. (6) and the consequent field equations. To make these models credible requires finding a source for the implied repulsive gravity. To attribute the repulsive gravity to a mysterious dark ‘energy’ represented by a cosmological constant Λ would be to beg the question and would add nothing to our understanding of the cosmos. As it happens there is ready to hand a well-aged theoretical discovery that has negative, repulsive gravity as an integral feature. In 1973 I described in considerable detail a model of a gravitating particle alternative to the Schwarzschild vacuum solution of Einstein’s field equations. This space-time manifold, which I termed a ‘drainhole’ with ‘ether’ flowing through it, was discovered independently at about the same time




FIG. 6: Postbounce graph of $\log_2 S(t)$ versus $\log_2(1 + \bar{t})$ ($\bar{t} := t \text{ s}^{-1}$) for the best-fit solution with $H_0 = 74.2 \text{ km s}^{-1}/\text{Mpc}$ and $k = 0$. The early stage rapid inflation, after producing 177 doublings of the normalized scale factor S in the first second after the bounce, gives way smoothly to a long period of uphill, decelerative ‘coasting’ (where the graph is nearly linear). Initially, the acceleration parameter $Q := (\ddot{R}/R)/(\dot{R}/R)^2 = (\ddot{S}/S)/(\dot{S}/S)^2$ is positive and huge ($Q(0) = \infty$). In the coasting period $Q(t) \approx Q_{\min} = -0.5 + 7.33 \times 10^{-98}$, which is reflected in the observation that in that interval $\log_2 S(t) \approx 177 + ((217 - 177)/(58 - 0)) \log_2(1 + \bar{t}) = 177 + (20/29) \log_2(1 + \bar{t}) \approx 177 + (2/3) \log_2(1 + \bar{t})$, so that $S(t) \approx 2^{177} (1 + \bar{t})^{2/3} \approx 2^{177} \bar{t}^{2/3}$ ($S(t) \propto \bar{t}^{2/3} \implies Q(t) \equiv -1/2$). After the coasting era $Q(t)$ becomes positive again at redshift $z_a = 0.684$ (when $t = 6.60$ Gyr and $\log_2(1 + \bar{t}) = 57.6$), rising to 0.557 at $t = t_0$, and then asymptotically to 1 as $t \rightarrow \infty$, the growth rate of $S(t)$ thus becoming asymptotically exponential. The corresponding graphs for the best-fit solutions with H_0 the same and $k = \pm 1$ are visually indistinguishable from this one.

by K. A. Bronnikov and formally some years earlier by O. Bergmann and R. Leipnik who rejected it for “physical reasons” [10–12]. It has come to be recognized as an early (apparently the earliest) example of what is now called by some a ‘traversable wormhole’, and has been analyzed from various perspectives by others [13–27]. The metric is a static, spherically symmetric solution of the field equations (7) and (8) with $\mu = \bar{\mu} = 0$ and $\psi = 0$. (NB $\mathbf{R}_{\alpha\beta}$ and $\mathbf{R}$ here are the negatives of those in [10].) It has the proper-time forms (in units in which $c = 1$)

$$d\tau^2 = [1 - f^2(\rho)] dT^2 - [1 - f^2(\rho)]^{-1} d\rho^2 - r^2(\rho) d\Omega^2 \quad (60)$$

$$= dt^2 - [d\rho - f(\rho) dt]^2 - r^2(\rho) d\Omega^2, \quad (61)$$

where $t = T - \int f(\rho) [1 - f^2(\rho)]^{-1} d\rho$,

$$f^2(\rho) = 1 - e^{-(2m/n)\alpha(\rho)}, \quad (62)$$

$$r(\rho) = \sqrt{(\rho - m)^2 + a^2} e^{(m/n)\alpha(\rho)}, \quad (63)$$

and

$$\phi = \alpha(\rho) = \frac{n}{a} \left[\frac{\pi}{2} - \tan^{-1} \left(\frac{\rho - m}{a} \right) \right], \quad (64)$$

with $a := \sqrt{n^2 - m^2}$, the parameters m and n satisfying $0 \leq m < n$. (The coordinate ρ used here translates to $\rho + m$ in [10].) The shapes and linear asymptotes of r and f^2 are shown in Fig. 7. Not obvious, but verifiable, is that $f^2(\rho) \sim 2m/\rho$ as $\rho \rightarrow \infty$, which, together with $r(\rho) \sim \rho$ as $\rho \rightarrow \infty$, shows m to correspond to the Schwarzschild mass parameter.

The shared metric of the cross sections of constant t is $d\rho^2 + r^2(\rho) d\Omega^2$. The choke point of the drainhole throat is the 2-sphere at $\rho = 2m$, of areal radius $r(2m) = ne^{(m/n)\alpha(2m)}$, which increases monotonically from n to ne as m increases from 0 to n . Thus the size of the throat

is determined primarily by n , and only secondarily by m . Although the scalar field ϕ has a nonkinetic ‘energy’ density that contributes to the space-time curvature through $T_{\alpha\beta}$, this energy has little to do with the strength of gravity (as determined by m), rather is associated with negative spatial curvatures found in the open throat, curvatures whose negativity mandates the unconventional choice for the polarity of the coupling of ϕ to the space-time geometry, as expressed by a minus sign in $T_{\alpha\beta}$ in Eq. (7). Even when $m = 0$, so that there is no gravity, these negative curvatures persist and the throat remains open. As a matter of perspective, it is more insightful to consider that the scalar field does not *cause* (i. e., is not a source of) these spatial curvatures, but simply tells of their existence and describes their configuration. This perspective helps disabuse one of the peculiar notion that geometrically unexceptionable space-time manifolds such as the drainhole are somehow a product of ‘exotic’ matter just because their Ricci tensors disrespect some ‘energy condition’. Moreover, it is not a great stretch to surmise that, whereas the parameter m specifies the active gravitational mass of the (nonexotic) drainhole particle, the size parameter n specifies in some way its inertial rest mass. This speculation is supported by two considerations: first, as shown in [10], the total energy of the scalar field ϕ lies in the interval from $n/2$ to $n\pi/2$, thus is essentially proportional to n ; second, it would seem likely that the bigger the hole, the greater the force needed to move it. A ‘higgsian’ way of expressing this idea is to say that the drainhole ‘acquires’ (inertial) mass from the scalar field ϕ .

Because $r(\rho) \geq n > 0$ and $f^2(\rho) < 1$, the drainhole space-time manifold is geodesically complete and has no one-way event horizon, the throat being therefore traversable by test particles and light in both directions. The manifold is asymptotic as $\rho \rightarrow \infty$ to a Schwarzschild manifold with (active gravitational) mass parameter m . The flowing ‘ether’ (a figurative term for a cloud of inertial observers free-falling geodesically from rest at $\rho = \infty$) has radial velocity $f(\rho)$ (taken as the negative square root of $f^2(\rho)$) and radial acceleration $(f^2/2)'(\rho)$, which computes to $-m/r^2(\rho)$ and therefore is strongest at $\rho = 2m$, where r assumes its minimum value. Because the radial acceleration is everywhere aimed in the direction of decreasing ρ , the drainhole attracts test particles on the high, front side, where $\rho > 2m$, and repels them on the low, back side, where $\rho < 2m$. Moreover, the manifold is asymptotic as $\rho \rightarrow -\infty$ to a Schwarzschild manifold with mass parameter $\bar{m} = -me^{m\pi/a}$, so the drainhole repels test particles more strongly on the low side than it attracts them on the high side, in the ratio $-\bar{m}/m = e^{m\pi/\sqrt{n^2-m^2}}$. The drainhole is a kind of natural accelerator of the ‘gravitational ether’, drawing it in on the high side and expelling it more forcefully on the low side. To replace the somewhat disreputable term ‘ether’ with something more acceptable in polite society




FIG. 7: Graphs of $r(\rho)$ and $f^2(\rho)$ for generic values of the parameters m and n ($0 \leq m < n$ and $a := \sqrt{n^2 - m^2}$).

one can imagine that it is space itself that is flowing into and through the drainhole. This substitution, which accords with Einstein’s insight that the concepts of space and of a gravitational ether are essentially interchangeable [28], should cause no alarm, for the very notion of an expanding universe already ascribes to space the requisite plasticity.

The discovery of the drainhole manifolds arose, in my case, from a search for a model for gravitating particles that, unlike a Schwarzschild space-time manifold, would have no singularity. Geodesic completeness and absence of event horizons resulted naturally from that requirement and the invoking of a minimally coupled scalar field to weaken the field equations. As shown in [10], a drainhole possesses all the geodesic properties that a Schwarzschild blackhole possesses other than those that depend on the existence of its horizon and its singularity, having evicted the horizon and replaced the singularity with a topological passageway to another region of space. Drainholes are able, therefore, to reproduce all the externally discernible aspects of physical blackholes (if such things exist) that Schwarzschild blackholes reproduce. That their back sides have never been knowingly observed (but in principle could be) is no more troubling

than the impossibility of directly observing the back sides of Schwarzschild horizons. For these reasons drainholes are more satisfactory than Schwarzschild blackholes as mathematical models of centers of gravitational attraction. Moreover, there is little reason to doubt that rotating drainhole manifolds analogous to the Kerr rotating blackhole manifolds exist and will prove to be better models than the Kerr manifolds. (A relatively recent solution of the field equations (7) and (8) perhaps describes such a manifold [26].)

B. Dark matter and dark ‘energy’ from drainholes

A physical center of attractive gravity modeled by a drainhole would qualify to be called a ‘darkhole’, inasmuch as (as shown in [10]) it would capture photons that venture too close, but, unlike a blackhole, must eventually release them, either back to the attracting high side whence they came or down through the drainhole and out into the repelling low side. Thus one can imagine that at galactic centers will be found not supermassive blackholes, but supermassive darkholes. At the center of our galaxy, for example, instead of a blackhole of Schwarzschild mass $m \approx 4 \times 10^6 M_\odot \approx 1.18 \times 10^7$ km (in geometric units), and of horizon area $4\pi(2m)^2 \approx 4.39 \times 10^{14}$ km², there might reside a drainhole the area $4\pi r^2(2m)$ of whose throat would lie between $4\pi n^2$ and $4\pi(ne)^2$, with n constrained only by having to exceed m . As measured by the bounds n and ne on the areal radius $r(2m)$, and the corresponding bounds $-m/4\pi n^2$ and $-m/4\pi(ne)^2$ on the maximum radial acceleration $(f^2/2)'(2m) = -m/r^2(2m)$, such an object might be large but relatively benign ($n \gg 0$, $m/n \ll 1$), large and not so benign ($n \gg 0$, $m/n \approx 1$), small and malignant ($n \approx 0$, $m/n \approx 1$), small and not so malignant ($n \approx 0$, $m/n \ll 1$), or something in between.

There is, however, more to the story. A central tenet of the general theory of relativity is that every elementary object that ‘has gravity’ is a manifestation of a local departure of the geometry of space-time from flatness. If such an object has other properties ascribed to it by quantum theory, these must be additional to the underlying geometric structure. I therefore propose the hypothesis that every such elementary gravitating object is at its core an actual physical drainhole — these objects to include not only elementary constituents of visible matter such as protons and neutrons, or, more elementarily, quarks, but also the unseen particles of ‘dark matter’ whose existence is at present only inferential. Moreover, I will consider that visible matter and primordial matter are, gravitationally, one and the same, and that the dark matter is the continuously created matter.

The pure, isolated drainhole described by Eqs. (60–64) is an ‘Einstein–Rosen bridge’ between two otherwise disjoint ‘subuniverses’, each of which, if evolving, would by itself consume for its description all the resources of a Robertson–Walker metric. Nonisolated drainholes could

presumably exist not only as ‘bridges’ between our subuniverse and another (or multiple others), but also as ‘tunnels’ from one place in our subuniverse to another, possibly quite distant from the first by every nontunnel route. Both types could contribute to the negative mass density $\bar{\mu}$ as well as to μ , the bridge drainholes contributing to $\bar{\mu}$ if their gravitationally repulsive back sides reside in our subuniverse, the tunnel drainholes contributing to $\bar{\mu}$ by way of their gravitationally repulsive exit portals. Tunnel drainholes are easy enough to visualize in abundance as topological holes into which space disappears, only to reappear elsewhere in our subuniverse, in analogy with rivers that go underground and surface somewhere downstream. Having both portals located in our subuniverse, tunnel drainholes would have properties we could fully take account of. Bridge drainholes, on the other hand, with only one side in our subuniverse would have properties dependent in part on circumstances in other subuniverses, circumstances beyond our ken. I shall, therefore, assume that no bridge drainholes contribute to $\bar{\mu}$ (or to μ), that the only contributors are tunnel drainholes, consequently that our subuniverse, even if part of a ‘multiverse’, is self-contained.

Lacking for the present a full mathematical description of these tunnel drainholes, let us nevertheless proceed as if they exist and are characterized by parameters m and n related as in an isolated bridge drainhole. Every particle of gravitating matter, whether in the ‘primordial’ (P) category or the ‘continuously created’ (C) category, is then at its core one of these tunnels, and it becomes a question of relating m and n to the densities $\mu_{P,0}$, $\bar{\mu}_{P,0}$, $\mu_{C,0}$, and $\bar{\mu}_{C,0}$.

The fitting of the cosmological solutions of Sec. IV to the SNe Ia data determines, by way of Eqs. (19), (20), (38), and (40), and Table II, the net densities $\mu_{P,0} + \bar{\mu}_{P,0}$ and $\mu_{C,0} + \bar{\mu}_{C,0}$, but tells nothing about the mix between the positive densities and the negative. To ‘fix the mix’, even if somewhat arbitrarily and only for P-densities, a procedure comes to mind that at first glance appears to replace a negative net density with a positive density, but on closer inspection is seen not do so. Equation (42) can be recast in the form

$$H^2 = \frac{c^2}{3} A_C - \frac{8\pi\kappa}{3} (\mu_{P,0} + \bar{\mu}_{P,0}) \frac{1}{\tilde{R}^3} + H_0^2 \left(\Omega_k \frac{1}{\tilde{R}^2} + \Omega_B \frac{1}{\tilde{R}^6} \right). \quad (65)$$

The analogous equation for the ‘concordance’ (Λ CDM) model is

$$H^2 = \frac{c^2}{3} \Lambda + \frac{8\pi\kappa}{3} \mu_M \frac{1}{\tilde{R}^3} + H_0^2 \left(\Omega_k \frac{1}{\tilde{R}^2} + \Omega_R \frac{1}{\tilde{R}^4} \right), \quad (66)$$

where the subscripts M and R refer to matter and radiation, respectively. As the only kind of gravitating matter

contemplated by the concordance model is effectively primordial matter (existent from the epoch of baryon genesis or earlier, treated in Eq. (66) as if existent from the big bang onward), the density μ_M can be identified with the density $\mu_{P,0}$, and then the corresponding primordial matter terms in the two equations will match exactly if $-(\mu_{P,0} + \bar{\mu}_{P,0}) = \mu_M = \mu_{P,0}$, thus if $\bar{\mu}_{P,0}/\mu_{P,0} = -2$.

To produce P-densities in the ratio $\bar{\mu}_P/\mu_P$ with tunnel particles of uniform masses m_P and sizes n_P requires that $\bar{\mu}_P/\mu_P = \bar{m}_P/m_P = -e^{m_P\pi/\sqrt{n_P^2 - m_P^2}}$, which entails that $\bar{\mu}_P/\mu_P$ is constant and

$$\left(\frac{m_P}{n_P}\right)^2 = \frac{[\ln(-\bar{\mu}_P/\mu_P)]^2}{[\ln(-\bar{\mu}_P/\mu_P)]^2 + \pi^2}. \quad (67)$$

If indeed $\bar{\mu}_{P,0}/\mu_{P,0} = -2$, then $\bar{\mu}_P/\mu_P = -2$, and $m_P/n_P = \ln 2/\sqrt{(\ln 2)^2 + \pi^2} \approx 0.22$. Under the assumption that n_P , or better, the throat radius $r(2m_P)$, is a proportional measure of the inertial mass of the primordial particles, this fixing of m_P/n_P would provide a basis outside the context of newtonian physics for suspecting the existence of a uniform ratio of active gravitational mass to passive-inertial mass of those particles. If, as I propose, the primordial particles are taken to be the gravitational cores of the protons and neutrons of the present epoch, then perhaps $r(2m_P) = m_{p-n}$, the inertial mass of a proton or a neutron. Converted to geometric units $m_{p-n} = 1.67 \times 10^{-24} \text{ g} = 1.24 \times 10^{-52} \text{ cm}$, so that $n_P = r(2m_P)e^{-(m_P/n_P)\alpha(2m_P)} = 1.24 \times 10^{-24} \text{ g} = 9.21 \times 10^{-53} \text{ cm}$ and $m_P = 2.67 \times 10^{-25} \text{ g} = 1.99 \times 10^{-53} \text{ cm}$. From Eqs. (19) and (40) one calculates that

$$\mu_{P,0} + \bar{\mu}_{P,0} = -\frac{3H_0^2}{8\pi\kappa} \Omega_P, \quad (68)$$

which for the best-fit value $\Omega_P = 0.295$ of Table II (cf. Table I) reduces to $\mu_{P,0} + \bar{\mu}_{P,0} = -3.05 \times 10^{-30} \text{ g/cm}^3 = -2.27 \times 10^{-58} \text{ cm/cm}^3$. With $\bar{\mu}_{P,0}/\mu_{P,0} = -2$ this gives $\mu_{P,0}/m_P = 11.4/\text{m}^3$ as the present proton-neutron (*qua* primordial particle) number density.

Comparison of the terms $c^2 A_C/3$ and $c^2 \Lambda/3$ in Eqs. (65) and (66) serves only to provide an identity for Λ , so to ‘fix the mix’ of positive and negative densities in $\mu_{C,0} + \bar{\mu}_{C,0}$ requires a strategy different from that used for $\mu_{P,0} + \bar{\mu}_{P,0}$. Note that, unlike the concordance model, which combines the gravitational effects of baryonic (visible) matter and cold dark matter in the term $(8\pi\kappa/3)(\mu_M/\tilde{R}^3)$ in Eq. (66) and thereby predicts from observational data a cold dark matter (CDM) to baryonic matter (BM) mass ratio of approximately 4:1, presumably constant in time, the model developed here reserves the analogous term in Eq. (65) for the effects of baryonic matter alone. That leaves the CDM to BM ratio, specified here as μ_C/μ_P , undetermined, and moreover allows it to be time-dependent, inasmuch as $\mu_C + \bar{\mu}_C$ is constant and $\mu_P + \bar{\mu}_P \propto R^{-3}$. To generate estimates for C-matter like those for P-matter, let us set $\gamma(z) := \mu_C(t)/\mu_P(t)$,

where t is the epoch of events seen at redshift z , and use $\gamma_0 := \gamma(0) = \mu_C(t_0)/\mu_P(t_0)$ as a parameter. From Eqs. (20) and (38) one calculates that

$$\mu_{C,0} + \bar{\mu}_{C,0} = -\frac{3H_0^2}{4\pi\kappa} \Omega_C. \quad (69)$$

This, together with Eq. (68) and the stipulation that $\bar{\mu}_{P,0}/\mu_{P,0} = -2$, implies that

$$\frac{\bar{\mu}_{C,0}}{\mu_{C,0}} = -\left(1 + \frac{2\Omega_C}{\gamma_0 \Omega_P}\right). \quad (70)$$

If, as in the case of P-particles, we assume that all the tunnel C-particles have the same mass parameter m_C and same size parameter n_C , then, just as for P-particles, we have that $\bar{\mu}_C/\mu_C$ is constant and

$$\left(\frac{m_C}{n_C}\right)^2 = \frac{[\ln(-\bar{\mu}_C/\mu_C)]^2}{[\ln(-\bar{\mu}_C/\mu_C)]^2 + \pi^2}. \quad (71)$$

Using this equation in conjunction with Eq. (70) and the equation $n_C = r(2m_C)e^{-(m_C/n_C)\alpha(2m_C)}$, we can construct a table of output values of parameters of interest for selected input values of γ_0 and $r(2m_C)$. The result is Table III, which uses $H_0 = 74.2 \text{ km s}^{-1}/\text{Mpc}$ and the best-fit values $\Omega_P = 0.295$ and $\Omega_C = 0.705$ from Table II (cf. Table I).

This discussion has assumed that the mass and size parameters of the P- and C-particles remain constant over time, which need not be the case. Indeed, it is conceivable, perhaps even likely, that these particles would share in the universal contraction and expansion. A particle that did so could be represented by a drainhole model that over the course of time from $-\infty$ to ∞ shrinks from infinite size to a point and then grows back to infinite size. Such a model is presented in [29] as a solution of the same field equations that the static drainholes satisfy, namely, Eqs. (7) and (8) with $\mu = \bar{\mu} = 0$ and $\psi = 0$. This model has $m = 0$, thus is devoid of gravity, but it could in principle be modified to one that has $m \neq 0$. Allowing the P- and C-particles to contract and expand with the universe would, of course, necessitate a revision of the discussion above and of Table III.

VI. ISSUES AND OBSERVATIONS

The conceptual basis of the model of the cosmos developed in the preceding sections differs in many respects from that of the concordance model. These differences and their consequences raise a number of issues, some of which have been discussed above to greater or lesser extent, others of which have been passed over. In this section I shall specify and comment on those that I have been able to identify along the way, in approximately the order in which they appeared. For ease of reference I shall call my model the PCB model, and the concordance model the Λ CDM model.

TABLE III: Computed parameter values for continuously created C-matter: μ_C (attractive mass density), m_C (attractive mass of a particle), $\bar{\mu}_C/\mu_C$ (ratio of repulsive mass density to attractive mass density), and μ_C/m_C (particle number density). Inputs are a selection of assumed values of $r(2m_C)$ (inertial mass of a particle) and γ_0 (ratio at present of attractive mass density of C-matter to that of primordial P-matter, proposed as dark matter to baryonic matter density ratio). The best-fit values $\Omega_P = 0.295$ and $\Omega_C = 0.705$ are taken from Table II (cf. Table I), with $H_0 = 74.2 \text{ km s}^{-1}/\text{Mpc}$.

Inputs		Outputs			
$r(2m_C)$	$\gamma_0 := \frac{\mu_{C,0}}{\mu_{P,0}}$	$\mu_C \text{ (g/cm}^3\text{)}$	$m_C \text{ (g)}$	$\frac{\bar{\mu}_C}{\mu_C}$	$\frac{\mu_C}{m_C} \text{ (1/m}^3\text{)}$
$m_{p-n} = 1.67 \times 10^{-24} \text{ g}$ $= 1.24 \times 10^{-52} \text{ cm}$	10	3.05×10^{-29}	1.72×10^{-25}	-1.48	1.77×10^2
	100	3.05×10^{-28}	2.43×10^{-26}	-1.05	1.26×10^4
	200	6.10×10^{-28}	1.24×10^{-26}	-1.02	4.91×10^4
	500	1.53×10^{-27}	5.04×10^{-27}	-1.01	3.03×10^5
$10^2 \times m_{p-n}$	10	3.05×10^{-29}	1.72×10^{-23}	-1.48	1.77
	100	3.05×10^{-28}	2.43×10^{-24}	-1.05	1.26×10^2
	200	6.10×10^{-28}	1.24×10^{-24}	-1.02	4.91×10^2
	500	1.53×10^{-27}	5.04×10^{-25}	-1.01	3.03×10^3
$10^{10} \times m_{p-n}$	10	3.05×10^{-29}	1.72×10^{-15}	-1.48	1.77×10^{-8}
	100	3.05×10^{-28}	2.43×10^{-16}	-1.05	1.26×10^{-6}
	200	6.10×10^{-28}	1.24×10^{-16}	-1.02	4.91×10^{-6}
	500	1.53×10^{-27}	5.04×10^{-17}	-1.01	3.03×10^{-5}
$m_{\text{Planck}} = 2.18 \times 10^{-5} \text{ g}$ $= 1.62 \times 10^{-33} \text{ cm}$ $= 1.31 \times 10^{19} \times m_{p-n}$	10	3.05×10^{-29}	2.24×10^{-6}	-1.48	1.36×10^{-17}
	100	3.05×10^{-28}	3.16×10^{-7}	-1.05	9.65×10^{-16}
	200	6.10×10^{-28}	1.62×10^{-7}	-1.02	3.77×10^{-15}
	500	1.53×10^{-27}	6.56×10^{-8}	-1.01	2.33×10^{-14}

A. Gravity and passive-inertial mass

As noted at the beginning, Newton's law of action and reaction allows the inference that the ratio m_a/m_{p-i} of active gravitational mass to passive-inertial mass is the same for all 'bodies', that is, for all matter in bulk. Because Newton's theory describes gravity as well as it does, this inference must be at least approximately correct. Even if exact, however, it is only a statement about a numerical ratio of quantities that *conceptually* have nothing to do with one another: the generation of a gravitational field on the one hand, the resistance to being accelerated by a field of any sort on the other. To try to force a marriage between these concepts is akin to the proverbial task of trying to push a square peg into a round hole. If one were to treat this unlikely union as legitimate, then

one would have to say that a gravitating body flying by at speed v would have its active gravitational mass and therefore its gravity increased by a factor $1/\sqrt{1-v^2/c^2}$ because that's what happens to its inertial mass. But the 'increase of inertial mass' by that factor has nothing to do with an innate property of the body such as the strength of its gravity, rather is simply a quantification of the fact that the closer any body's speed is to the speed of light, the harder it is to accelerate that body with a field that propagates at the speed of light. (The analogy of a person on foot pushing a dead-battery car to get it started comes to mind: if the car can't be started at a speed less than the pusher's maximum running speed, it won't be started.) And let us note in passing that when an electrically charged particle's electromagnetic field is observed to depend on the relative velocity of the observer and the particle, this is attributed to the fact that

the field propagates at finite speed, not to a change in the particle's electric charge.

If passive-inertial mass is not a source of gravity, then as said in Sec. II Einstein's 'energy-tensor of matter' $T^{\alpha\beta} := \mu u^\alpha u^\beta + (p/c^2)(u^\alpha u^\beta - g^{\alpha\beta})$ has no role to play in the field equations of gravity, so one loses the spuriously attractive feature that the (Einstein) field equations imply that $T_{\alpha\beta;\beta} = 0$, interpreted as the equations of motion of the gravitational source matter. This, however, does not keep one from asserting that $T_{\alpha\beta;\beta} = 0$; it only requires that one look elsewhere for its justification. But in making that assertion one must recognize that, for $T_{\alpha\beta;\beta} = 0$ to be so interpreted, μ must be the density of inertial-passive mass, not of active gravitational mass.

All too frequently there appear in the popular press accounts of astronomers 'weighing' the universe or 'weighing' a galaxy. One can object, of course, to the presumably intentional blurring of the distinction between inertial mass and weight, but a larger objection is that such accounts leave the reader with the false impression that determining an entity's active gravitational mass is known to be equivalent to determining its passive-inertial mass (and therefore its weight). Only if one can prove the existence of a universal proportionality between the two can one legitimately claim to have 'weighed' the universe or a galaxy, and even then only indirectly. Unless and until there is such a proof, pretending that the universe or a galaxy has been weighed spreads ignorance, not knowledge. The most elementary unit of knowledge is a distinction made — a fact built in to every digital computer. Conversely, the most elementary unit of ignorance is a distinction not made.

B. Choice of a variational principle for gravity

The step up in Sec. II from the nonrelativistic variational principle $\delta \int (|\nabla \phi|^2 + 8\pi\kappa\mu\phi) d^3x = 0$ that generates the Poisson equation $\nabla^2 \phi = 4\pi\kappa\mu$ to the relativistic principle $\delta \int (\mathbf{R} - \frac{8\pi\kappa}{c^2}\mu) |g|^{\frac{1}{2}} d^4x = 0$ is an application of Occam's Razor, and as such can only be justified retrospectively, by its consequences. Alone, this principle has, through its Euler-Lagrange equations, little to say about cosmology, but when modified by inclusion of the cosmological constant Λ it says something significant, namely, that there is gravitationally repulsive matter in the universe and there's more of it than there is of attractive matter. This is a consequence that has consequences, but to get them all requires the next step up, to the principle of Eq. (6) which has in it in addition to the positive and negative mass densities μ and $\bar{\mu}$ the scalar fields ϕ and ψ .

One will recall that in deriving the field equations (7) and (8) only the metric components and not the densities or the scalar fields were varied in Eq. (6). Not varying the densities is normal, but not varying the scalar fields goes against orthodox practice. The rationale for leaving

the scalar fields unvaried is this: In a space-time manifold everything is determined by the metric, and that should be all there is to be fixed. To vary the scalar fields would be to treat them as something extraneous to the metric, whereas their proper role should be to help describe the metric, not to serve as a source of some sort.

Looked at another way, including the terms $-\frac{8\pi\kappa}{c^2}\mu$, $-\frac{8\pi\kappa}{c^2}\bar{\mu}$, $2\phi^\gamma\phi_{,\gamma}$, and $-2\psi^\gamma\psi_{,\gamma}$ in the action integrand can be seen as accounting for internal constraints on the metric, analogously to the way including a Lagrange multiplier λ (Λ in the Λ CDM case) accounts for an external constraint. Just as one does not vary λ (or Λ), one should not vary μ , $\bar{\mu}$, ϕ , or ψ . In this role μ , $\bar{\mu}$, ϕ , and ψ simply relax the field equations to allow a larger class of metrics to satisfy them. The relaxed field equations are as useful as the metrics that satisfy them, no more, no less.

The inclusion of the scalar field ψ coupled to the metric with polarity opposite to that of ϕ 's coupling is dictated in the first instance by the observation that if the β^2 term were absent from Eq. (21), then, once $R(t)$ surpassed $\sqrt[3]{B/A_P}$ (which is $R_{H_{\max}}$ when $k = 0$), Eq. (21) would be false. Moreover, if the shoe were on the other foot and the α^2 term were absent, then Eq. (21) would be false when $R(t)$ was less than $\sqrt[3]{B/A_P}$. Thus both ϕ and ψ are needed, coupled to the geometry with opposite polarities. That the combination $2\phi^\gamma\phi_{,\gamma} - 2\psi^\gamma\psi_{,\gamma}$ in the variational principle of Eq. (6) is the real part of $2\chi^\gamma\chi_{,\gamma}$, where $\chi := \phi + i\psi$, suggests that space-time as seen here might be a restriction of a more general, at least partially complexified space-time geometry.

In the drainhole model described in [10] and here in Sec. V only the scalar field ϕ was used, and it was assumed to satisfy the field equation $\square\phi = 0$, obtained in [10] by varying ϕ in the variational principle. That varying, however, ought to have been omitted, as $\square\phi = 0$ was already a consequence of the equations obtained from varying only the metric. In the drainhole model, therefore, ϕ is not a source of anything, rather is just a descriptor of the geometry of space-time — and primarily, as seen by setting $m = 0$ in Eqs. (62) and (63), of space alone. That the polarity of its coupling to geometry is opposite to the orthodox is precisely what allows the negative curvatures necessary to the existence of the open drainhole throat. To say that a scalar field coupled to the space-time geometry with this polarity is a 'ghost' or 'phantom' field associated with 'exotic matter' is simply to indulge in misleading jargon, an error whose root can be traced to an uncritical acceptance of Einstein's assumption that energy produces gravity.

C. Inflation and the 'big bounce'

If the field equations dictate that R have a positive minimum value $R_{\min}$, then the history of the universe will include a bounce off that minimum; if not, then in-

stead of a bounce there will be a ‘bang’, where $R = 0$. As is evident from the definition of $P_1(R)$ in Eq. (24), as well as from examining Fig. 1, for $R_{\min}$ to be positive it is necessary and sufficient that the integration constant B be positive (otherwise $P_1(R) = -B \geq 0$ at $R = 0$, which would make $R_{\min} = 0$). Also evident is that the smaller a positive B is, the closer $R_{\min}$ is to 0; in fact, as noted in Sec. III, $R_{\min} \sim \sqrt[3]{B/2A_P}$ as $B \rightarrow 0$. Moreover, it is clear that the term $2B/3R^6$ in Eq. (23) is the term that produces rapid inflation when R is small by making $\ddot{R}$ large, but only when $B > 0$. And because $B - A_P R^3 \sim B - A_P(B/2A_P) = B/2$ as $B \rightarrow 0$, positivity of B demands that the $\dot{\alpha}^2$ term be present in Eq. (21), thus that the scalar field ϕ be present in the field equations, coupled to the space-time geometry with the unconventional polarity. This is the way in which relaxing the field equations by including ϕ allows the geometry to produce both a ‘bounce’ and the inflation that turns it into a ‘big bounce’. Let us note that this is accomplished without invoking a ‘slow roll down a potential’, a common element in many proposed inflationary scenarios. Indeed, that ϕ is treated merely as an aid in describing the space-time geometry rather than as a ‘physical source’ of the geometry makes the notion of introducing a potential function for ϕ irrelevant.

D. The PCB model vis-à-vis the Λ CDM model

After the matching $-(\mu_{P,0} + \bar{\mu}_{P,0}) = \mu_M = \mu_{P,0}$ in Sec. V the only functional difference between the expressions for H^2 in the PCB model and in the Λ CDM model lies in their terms of highest order in $1/\tilde{R}$: $H_0^2 \Omega_B/\tilde{R}^6$ in Eq. (65) and $H_0^2 \Omega_R/\tilde{R}^4$ in Eq. (66). In fully normalized form these equations read

$$H^2 = H_0^2 \left(\Omega_C + \Omega_k \frac{1}{\tilde{R}^2} + \Omega_P \frac{1}{\tilde{R}^3} + \Omega_B \frac{1}{\tilde{R}^6} \right) \quad (72)$$

(which is Eq. (42)) and

$$H^2 = H_0^2 \left(\Omega_\Lambda + \Omega_k \frac{1}{\tilde{R}^2} + \Omega_M \frac{1}{\tilde{R}^3} + \Omega_R \frac{1}{\tilde{R}^4} \right). \quad (73)$$

Absent their highest-order terms, and with the obvious identifications among the remaining Ω s, these equations would produce, by way of the corresponding formulas for the distance modulus μ , exactly the same best fits to the SNe Ia data, the parameters and the χ^2 statistic being those of Table I. The rationale behind the construction of Table II was that, because the SNe Ia data extend into the past only to $z = 1.755$, restoring the Ω_B term in Eq. (72) while retaining the $\Omega_B = 0$ best-fit values of the other parameters will yield a fit with negligibly different χ^2 . The same reasoning could be expected to apply also to restoring the Ω_R term in Eq. (73). This I have tested,

with the result that so long as $\Omega_R \lesssim 10^{-5}$ the values computed for χ^2 will match those of Table I to the precision shown, larger values of Ω_R generating larger values of χ^2 (and younger ages for the universe). It is therefore the case that the PCB model is at least as consistent with the SNe Ia data as the Λ CDM model is.

The omission from the PCB-model field equations of a radiation term proportional to $1/R^4$ goes against conventional wisdom but is not without motivation. Inclusion of such a term would seem to violate the basic working hypothesis that energy is not a generator of gravity. Radiant energy, however, although convertible to kinetic energy through its influence on the motion of electrically charged objects, is not kinetic energy and therefore might be a gravity generator. This would presumably come about through a coupling of the electromagnetic field components $F_{\alpha\beta}$ to the space-time geometry. The difficulty is that the usual method of coupling by inserting a term proportional to $F^{\gamma\delta}F_{\delta\gamma}$ into the integrand of the variational principle has no geometrical justification and produces field equations of the sort that Einstein likened to a building with one wing made of fine marble ($\mathbf{R}_{\alpha\beta} - \frac{1}{2}\mathbf{R}g_{\alpha\beta}$), the other, of low grade wood (e. g., the energy-momentum tensor for F) [30]. What is needed is a version of the PCB model developed in the context of a geometrical unified field theory of the sort that Einstein spent his later years in an unsuccessful quest for. Although I have been able to construct such a theory as a hybrid of Weyl’s conformal theory and Kaluza’s five-dimensional theory [31, 32], the manner in which that theory’s variational principle and field equations connect the electromagnetic field to geometry does not suggest a way to relate F to the space-time geometry in the present context. For this reason I have chosen to omit the radiation term from the field equations governing the PCB model. Such a term can, of course, be reinstated on an *ad hoc* basis if a sufficiently compelling reason for doing so presents itself.

E. Dark matter, dark ‘energy’, and the ‘Cosmological Constant Problem’

In the Λ CDM model dark matter is lumped with baryonic matter in Ω_M as primordial matter whose density is proportional to $1/R^3$. The unseen, unknown stuff in Ω_Λ , whose density has no R dependence, is called dark ‘energy’ in deference to Einstein and his assumption that energy is a source of gravity. In the PCB model only baryonic matter is treated as belonging to the primordial Ω_P sector with $1/R^3$ density dependence, dark matter being shifted to the Ω_C sector with density kept constant by continuous creation of the dark C-matter. And there, instead of dark ‘energy’ represented by a cosmological constant Λ many orders of magnitude too small to be consistent with various origins proposed for this energy,

one finds the gravitationally attractive dark matter particles accompanied by their gravitationally repulsive back sides, their net effect represented by the constant A_C of Eq. (20), functionally equivalent to Λ and of the requisite smallness. This interpretation of ‘dark energy’ as the back side of dark matter provides a self-consistent solution of the so-called ‘Cosmological Constant Problem’.

F. Ratio of dark matter to baryonic matter

Because in the Λ CDM model the densities of dark matter and baryonic matter are both proportional to $1/R^3$, the ratio of their densities is constant in time. Estimates of that ratio obtained from observations of visible structures such as galaxies and galactic clusters and superclusters as they exist at the present epoch in the vicinity of our galaxy are therefore used in studies of the formation of these structures in the distant past. In the PCB model, on the other hand, the dark matter density is held constant through continuous creation of C-particles, so the ratio μ_C/μ_P of dark C-matter density to baryonic P-matter density is proportional to R^3 . Consequently, the value of this ratio at the epoch corresponding to redshift z is $\gamma(z) = \gamma_0/(1+z)^3$, where γ_0 is the ratio at present, several possible values of which are included in Table III for comparison purposes. Taking this dependence of the ratio on redshift as real would no doubt influence the outcomes of studies of structure formation.

G. Continuous creation of C-matter tunnels

The stipulation that the net density $\mu_C + \bar{\mu}_C$ of C-matter stay fixed while the universe is expanding requires that the drainhole tunnels considered to be the particles of C-matter come continuously into existence. By what mechanism might this happen? Wheeler’s notion of a ‘quantum foam’ of wormholes popping into and out of existence would not provide a satisfactory explanation, for to keep the density constant each newly created tunnel must either remain in existence or else upon dying be multiply replaced by new ones. A more useful idea is that the tunnels arise from the stretching of space as the universe expands. Once started, this would be a regenerative process, each new tunnel causing more stretching that would generate additional tunnels. Such a process, moreover, could be expected to produce particles in a given region of space at a rate proportional to the rate of increase of the volume of that region, thus maintaining a constant particle density. If, as was assumed in Sec. V, these particles are all created with the same mass and size parameters m_C and n_C , the net density would be held constant. As to how the process could begin, one looks to the primordial P-matter tunnels, presumed to have always been present, pushing the universe toward

expansion. In the postbounce era the C-particles would add their contribution, the ultimate result being an acceleration of the expansion. In the prebounce era the C-particles, abundant in the distant past, would be dying out with the shrinking of space as the universe contracted into the bounce.

If the primordial P-matter has ‘always’ existed, then the question of what ‘caused’ it to exist is better left to the philosophers. Conceivably, however, the P-matter tunnels came into existence as dark matter particles the same way the C-matter tunnels did, but in the inflationary space-stretching era after the bounce. Only later would their entrance portals have acquired the decorations that turned them into baryonic matter. The really fundamental unanswered question, though, is how did space and time come to exist, for, time excepted, all the elements of the PCB model — the tunnels and the scalar fields — are merely features of the geometry of space. In fact, the question can be reduced to the existence of space alone, for the Kaluza–Weyl theory described in [31, 32] is based on a construction that produces time (thus space-time) from three-dimensional space and, repeated, produces a secondary time (thus space-time–time) from space-time. And in the space-time stage of that theory there occurs naturally a drainhole model in an expanding universe which repels on the low side more strongly that it attracts on the high side and needs no auxiliary scalar field for its description [33].

That all the C-particles are created with the same mass and size parameters is only a convenient assumption. If this assumption is relaxed, then the densities μ_C and $\bar{\mu}_C$ might vary, in which case the proportionality of $\mu_C + \bar{\mu}_C$ to $1/R^3$ could cease to be exact. This variability could then be used in an *ad hoc* way to produce some of the putative behaviors that proposed modifications of the Λ CDM model put forth under the rubric ‘quintessence’ are intended to allow.

H. Voids, walls, filaments, and nodes

The PCB model, just as the Λ CDM model, treats matter as uniformly distributed in space, an apparently reasonable assumption on a large enough scale. Observationally, however, a major fraction of space appears to comprise voids nearly empty of matter, bordered by walls, filaments, and nodes in which resides most of the matter, both the dark and the visible (baryonic). The formation of this cellular structure is generally considered to be an evolutionary product of early small fluctuations in the density of primordial matter. Explanations along that line are attempts to realize what Peebles has called his “perhaps desperate idea . . . that the voids have been emptied by the growth of holes in the mass distribution” [34]. If besides matter that attracts gravitationally there is also repulsive matter, then that idea can be rein-

terpreted and the desperation perhaps reversed, by the simple observation that, whereas attractive matter wants to congregate, repulsive matter is reclusive, pushing all matter, attractive or repulsive, away from itself.

In the PCB model each tunnel particle would presumably be created with its entrance and its exit close to one another in the ambient space (if far apart, all the better). In the ambient space the entrance would attract the exit, but the exit would repel the entrance more strongly, so the two portals would drift apart. Apply this to a multitude of such particles and you would expect to see the exits spread themselves over regions from which they had expelled the entrances, regions therefore devoid of ordinary (i. e., attractive) matter. The entrances, on the other hand, being brought together by both their mutual attractions and the repulsion from the exits, would aggregate into walls, filaments, and nodes on the boundaries of the void regions, just as observed in the real universe. What is more, the walls, filaments, and nodes so created would likely be, in agreement with observation, more compacted than they would have been if formed by gravitational attraction alone, for the repulsive matter in the voids would increase the compaction by pushing in on the clumps of attractive matter from many directions with a nonkinetic, positive pressure *produced by* repulsive gravity, a pressure not to be confused with the negative pseudo-pressure conjectured in the confines of Einstein's assumption to be a *producer* of repulsive gravity. In this way the PCB model explains qualitatively the observed cellular structure of the universe with minimal or no reliance on fluctuations in the primordial matter density.

The validity of this explanation of the cellular structure could be tested independently of the tunnel particle hypothesis by numerical simulation of evolution of an expanding universe with both attractive matter and repulsive matter distributed throughout. An interesting topic for investigation would be the propagation of light through the voids, particularly as the repulsive matter in a void might tend to distribute itself nonuniformly.

I. Geometric units and inertial mass

In Sec. V, in order to populate Table III, the radius $r(2m_C)$ of the C-particle throats was set equal to various multiples of the proton-neutron inertial mass m_{p-n} , converted to geometric units to give the radius its requisite units of length. Because the conversion from inertial mass units to geometric units ($1g \rightarrow G/c^2 \times 1g \approx 7.426 \times 10^{-29}$ cm) involves Newton's gravitational constant G , such a conversion would seem to be at odds with the assumption that inertial mass does not generate gravity. The resolution of this apparent inconsistency, dictated by the proposition that everything that exists is one or another aspect of the geometry of the universe, is that the gram is an artificial unit of inertial mass used for convenience, and that the proper unit of inertial mass

is a length, 7.426×10^{-29} cm, for example. That necessitates that $G/c^2 = 1$. The 'geometry is all' proposition dictates also that the second must be considered an artificial unit standing in for a length unit. Taking that unit to be the light-second makes $c = 1$ and, consequently, $G = 1$, which equations are commonly taken as the defining equations of geometric units. Considering the proper units of inertial mass and time to be units of length does not, of course, prevent the use of the gram and the second in their customary contexts.

J. Galactic nuclei as drainholes

If instead of a blackhole at the center of our galaxy there is a drainhole, what might it look like? From afar the behavior of matter and radiation around such a drainhole would not differ in appearance from the behavior of matter and radiation around a blackhole of the same mass and approximately the same size. For both a Schwarzschild blackhole of mass m and a drainhole of mass m the radial equation of motion of a test particle is

$$\frac{d^2\rho}{d\tau^2} = -\frac{m}{r^2(\rho)} + (\rho - 3m)\left(\frac{d\Omega}{d\tau}\right)^2, \quad (74)$$

with $r(\rho)$ given by Eq. (63) for the drainhole and $r(\rho) = \rho$ for the blackhole. In both cases τ is the proper time of the test particle. Equation (74) shows that for the drainhole as well as for the blackhole $3m$ is its 'star-grabbing' radius, as any star whose orbit takes it down to $\rho = 3m$ or below will never again have $d\rho/d\tau > 0$. The shared metric of the cross sections of constant t being $d\rho^2 + r^2(\rho) d\Omega^2$, the cross sections of the blackhole are euclidean, so the geodesic radius $3m$ of the sphere from which no star can escape is also its areal radius. For the drainhole the situation is different: the areal radius is $r(3m)$, which by increasing the size parameter n can be made as large as you like.

For purely radial motion the acceleration $-m/r^2(\rho)$ in Eq. (74) is the (nonvectorial) gradient $(f^2/2)'(\rho)$ of the specific kinetic energy $\frac{1}{2}v^2$ of observers free-falling from rest at $\rho = \infty$. It attains its maximum strength $m/r^2(2m)$ at the choke point of the drainhole throat. A measure of the tidal effects in the radial direction is the tidal gradient, given by

$$s(\rho) := \left(\frac{f^2}{2}\right)''(\rho) = \frac{2(\rho - 2m)}{(\rho - m)^2 + n^2 - m^2} \cdot \frac{m}{r^2(\rho)}, \quad (75)$$

which vanishes at the choke point and reaches a positive, stretching maximum

$$s_{\max} = \frac{\sqrt{3}m}{2n + \sqrt{3}m} \cdot \frac{1}{r^2(\rho_+)} \quad \text{at} \quad \rho_+ = 2m + \frac{n}{\sqrt{3}}, \quad (76)$$

and a negative, compressing minimum

$$s_{\min} = -\frac{\sqrt{3}m}{2n - \sqrt{3}m} \cdot \frac{1}{r^2(\rho_-)} \quad \text{at} \quad \rho_- = 2m - \frac{n}{\sqrt{3}}. \quad (77)$$

For a fixed m , $s_{\max}$ is a monotonic function of n , decreasing from

$$\lim_{n \rightarrow m} s_{\max} = \frac{3\sqrt{3}(2 - \sqrt{3})^2 e^{-(3-\sqrt{3})}}{2m^2} \quad (78)$$

to 0, while $s_{\min}$ increases monotonically from

$$\lim_{n \rightarrow m} s_{\min} = -\frac{3(12 + 7\sqrt{3})e^{-(3+\sqrt{3})}}{2m^2} \quad (79)$$

to 0, as n increases from m to ∞ .

Observations of stars near the galactic center put the central supermass constraining their motions at $m_{\text{gc}} \approx 4.31 \times 10^6 M_{\odot} = 6.36 \times 10^6 \text{ km}$ [35]. In ordinary units $1/m_{\text{gc}} = c^2/(6.36 \times 10^6 \text{ km}) = 1.41 \times 10^4 \text{ km/s}^2$ and $1/m_{\text{gc}}^2 = c^2/(6.36 \times 10^6 \text{ km})^2 = 2.22 \times 10^{-3} (\text{km/s}^2)/\text{km}$. Building on these, with the ratio n/m_{gc} as an input parameter, Table IV exhibits the corresponding values of the drainhole areal radius $r(2m)$, the star capture areal radius $r(3m)$, the maximum radial acceleration $m/r^2(2m)$, and the tidal gradients $s_{\max}$ and $s_{\min}$.

Stars, gases, and light falling into a supermassive blackhole encounter unbounded tidal stretching that is said to strip them of their identities as they approach and reach the central singularity, never to be seen again. Falling into a supermassive drainhole gases and light (and even stars if the ratio n/m_{gc} is large) can retain their identities and exit the drainhole on the other, gravitationally repelling side. This brings the following question: If supermassive drainholes at galactic centers are commonplace, and their exits reside in our (part of the) universe, where should we expect to see those exits, and how might we identify them? In accordance with the notion that the exits of drainhole tunnels tend to isolate themselves by pushing all other matter away, the obvious places to look would be the great voids filled with the exits of dark C-matter tunnel. Presumably they would appear as compact objects surrounded by a halo of outflowing gas, their light intrinsically blueshifted from its journey downward in the drainholes' gravitational potentials. The brightnesses of such objects would depend on the rate at which the gases and light were sucked into the drainhole entrances. If very bright, they might appear as isolated quasars. If dim, they might be present in the voids but unseen.

With a little imagination one can see how a supermassive drainhole might arise from a plenitude of baryonic P-matter and dark C-matter drainhole tunnels with their entrances brought tightly together by their mutual attractions. Begin with a pair of them, T_1 and T_2 say. If

not only their entrances but also their exits were close together, then the coalescing of T_1 and T_2 into a single tunnel might be feasible. If, however, as I have proposed, their exits had receded far from one another, then the only way T_1 and T_2 might reasonably coalesce into one tunnel would be to join at their entrances and close off into a single tunnel connecting their original exits. If that occurred, then the 'gravitational ether' flowing into their entrances would be diverted to the nearby entrances of one or more tunnels, thereby increasing their masses, both the attractive and the repulsive. The remnant of T_1 and T_2 , no longer connected to the ether flow, would soon become a sterile, gravityless tunnel connecting two distant places out in the voids. After many repetitions of this process there would be left a single tunnel large enough to accommodate all the combined ether flow of the original P- and C-matter tunnels — a supermassive drainhole.

One can equally well imagine that all the tunnels associated with baryonic matter and dark matter concentrated at a galactic center would retain their identities and their gravity, both the attractive and the repulsive, no matter how tightly packed or rapidly moving their entrances might be. At the tunnel exits there would be nothing new to look for, all the attractive gravity concentrated at their entrances being dispersed in the voids as negative gravity in the same way as it was before the concentration took place. Here there would be no merging of tunnels, thus no altering of the topology of space. Left to be analyzed mathematically would be only the simpler changes of spatial topology associated with the creation and destruction of the individual drainhole tunnels.

VII. SUMMARY

Cosmic inflation, deceleration, acceleration, dark matter, dark 'energy', and a 'big bounce', not to mention coasting, voids, walls, filaments, and nodes, have been found wrapped in one coherent package, namely, the variational principle $\delta \int [\mathbf{R} - \frac{8\pi\kappa}{c^2}(\mu + \bar{\mu}) + 2\phi^{\cdot\gamma}\phi_{\cdot\gamma} - 2\psi^{\cdot\gamma}\psi_{\cdot\gamma}] |g|^{\frac{1}{2}} d^4x = 0$ and the field equations it implies. Arriving at this variational principle depended on giving up Einstein's assumption that inertial mass and energy produce gravity.

VIII. NOTE

This is a revision and major extension of the earlier version(s) [36], retitled to reflect the revision.

TABLE IV: For a drainhole of mass parameter $m = m_{\text{gc}}$ and size parameter n , where $m_{\text{gc}} \approx 4.31 \times 10^6 M_{\odot} = 6.36 \times 10^6 \text{ km}$ in geometric units (the current best estimate of the supermass at the center of our galaxy), and a range of values of the ratio n/m_{gc} , this table exhibits the corresponding values of the drainhole areal radius $r(2m)$, the star capture areal radius $r(3m)$, the maximum radial acceleration $m/r^2(2m)$, and the maximum and minimum tidal gradients s_{max} and s_{min} .

Inputs	Outputs				
$\frac{n}{m_{\text{gc}}}$	$r(2m)$ (km)	$r(3m)$ (km)	$\frac{m}{r^2(2m)}$ (km s ⁻²)	s_{max} (km s ⁻² /km)	s_{min} (km s ⁻² /km)
1	1.73×10^7	2.10×10^7	1.91×10^3	1.17×10^{-4}	-7.08×10^{-4}
10	7.37×10^7	7.41×10^7	1.05×10^2	1.01×10^{-6}	-1.15×10^{-6}
100	6.46×10^8	6.46×10^8	1.37	1.39×10^{-9}	-1.41×10^{-9}
1000	6.37×10^9	6.37×10^9	1.41×10^{-2}	1.44×10^{-12}	-1.44×10^{-12}
10000	6.36×10^{10}	6.36×10^{10}	1.41×10^{-4}	1.44×10^{-15}	-1.44×10^{-15}
100000	6.36×10^{11}	6.36×10^{11}	1.41×10^{-6}	1.44×10^{-18}	-1.44×10^{-18}

- [1] A. Einstein, Die Grundlage der allgemeinen Relativitätstheorie, *Ann. der Physik* **49** (1916), 769–822, translated in *The Principle of Relativity* (Dover, New York, 1952), pp. 109–164.
- [2] G. Galilei, **Dialogues Concerning Two New Sciences** (Prometheus Books, Buffalo, N.Y., 1991), pp. 63 *et seq.*; http://galileoandeinstein.physics.virginia.edu/tns_draft/index.html, pp. 63 *et seq.*
- [3] A. Einstein, **Relativity, The Special and the General Theory: A Popular Exposition** (Methuen, London, 1916), (Bonanza Books, New York, 1961), pp. 66 *et seq.*; also available as an eBook (Routledge, London, New York, 2002).
- [4] D. F. Bartlett and D. Van Buren, Equivalence of active and passive gravitational mass using the moon, *Phys. Rev. Lett.* **57** (1986), 21–24.
- [5] L. B. Kreuzer, Experimental measurement of the equivalence of active and passive gravitational mass, *Phys. Rev.* **169** (1968), 1007–1012.
- [6] C. S. Unnikrishnan and G. T. Gillies, Do leptons generate gravity? First laboratory constraints obtained from some G experiments and possibility of a new decisive constraint, *Phys. Lett. A* **288** (2001), 161–166.
- [7] D. Hilbert, Die Grundlagen der Physik (Erste Mitteilung), *Nachr. Königl. Gesellschaft d. Wiss. Göttingen, Math.-Phys. Kl.*, 395–407; reprised in *Math. Annalen* **92** (1924), 1–32.
- [8] A. Riess, et al., New Hubble Space Telescope discoveries of Type Ia supernovae at $z > 1$: narrowing constraints on the early behavior of dark energy, *Astrophys. J.* **659** (2007), 98–121.
- [9] A. Riess, et al., A Redetermination of the Hubble Constant with the Hubble Space Telescope from a Differential Distance Ladder, *Astrophys. J.* **699** (2009), 539–563.
- [10] H. G. Ellis, Ether flow through a drainhole: a particle model in general relativity, *J. Math. Phys.* **14** (1973), 104–118; Errata: **15** (1974), 520.
- [11] K. A. Bronnikov, Scalar-tensor theory and scalar charge, *Acta Phys. Pol.* **B4** (1973), 251–266.
- [12] O. Bergmann and R. Leipnik, Space-time structure of a static spherically symmetric scalar field, *Phys. Rev.* **107** (1957), 1157–1161.
- [13] G. Clément, The Ellis Geometry (Letter to the editor), *Am. J. Phys.* **57** (1989), 967.
- [14] M. S. Morris and K. S. Thorne, Wormholes in spacetime and their use for interstellar travel: a tool for teaching general relativity, *Am. J. Phys.* **56** (1988), 395–412.
- [15] L. Chetouani and G. Clément, Geometrical-optics in the Ellis geometry, *Gen. Relativ. Gravit.* **16** (1984), 111–119.
- [16] G. Clément, Scattering of Klein-Gordon and Maxwell waves by an Ellis geometry, *Int. J. Theor. Phys.* **23** (1984), 335–350.
- [17] S. Kar, D. Sahdev, and B. Bhawal, Scalar waves in a wormhole geometry, *Phys. Rev. D* **49** (1994), 853–861.
- [18] C. Armendáriz-Picón, On a class of stable, traversable Lorentzian wormholes in classical general relativity, *Phys. Rev. D* **65** (2002), 104010–1–10.
- [19] H. Shinkai and S. A. Hayward, Fate of the first traversible wormhole: Black-hole collapse or inflationary expansion, *Phys. Rev. D* **66** (2002), 044005–1–9.
- [20] V. Perlick, Exact gravitational lens equation in spherically symmetric and static spacetimes, *Phys. Rev. D* **69** (2004), 064017–1–10.
- [21] A. Das and S. Kar, The Ellis wormhole with ‘tachyon matter’, *Class. Quantum Grav.* **22** (2005), 3045–3053.
- [22] K. K. Nandi, Y-Z. Zhang, and A. V. Zakharov, Gravitational lensing by wormholes, *Phys. Rev. D* **74** (2006), 024020–1–13.

- [23] T. Müller, Exact geometric optics in a Morris-Thorne wormhole spacetime, *Phys. Rev. D* **77** (2008), 044043-1–11.
- [24] D. K. Dey and S. Sen, Gravitational lensing by wormholes, *Mod. Phys. Lett. A* **23** (2008), 953–962.
- [25] A. G. Doroshkevich, N. S. Kardashev, D. I. Novikov, and I. D. Novikov, The Passage of Radiation through a Wormhole, *Astron. Rep.* **52** (2008), 685–691.
- [26] T. Matos and D. Nuñez, Rotating scalar field wormhole, *Class. Quantum Grav.* **23** (2006), 4485–4495.
- [27] K. K. Nandi, I. Nigmatzyanov, and R. Izmailov, New features of extended wormhole solutions in the scalar field gravity theories, *Class. Quantum Grav.* **25** (2008), 165020-1–19.
- [28] A. Einstein, Relativity and the ether, in *Essays in Science* (Philosophical Library, New York, 1934), pp. 98–111.
- [29] H. G. Ellis, The evolving, flowless drainhole: a nongravitating particle model in general relativity theory, *Gen. Relativ. Gravit.* **10** (1979), 105–123.
- [30] A. Einstein, Physics and reality, in *Essays in Physics* (Philosophical Library, New York, 1950), p. 39.
- [31] H. G. Ellis, Space-time-time: Five-dimensional Kaluza–Weyl space, [gr-qc/0107023](#).
- [32] H. G. Ellis, Space-time-time, [gr-qc/0205029](#).
- [33] H. G. Ellis, Darkholes: Nicer than blackholes — with a bright side, too (Does energy produce gravity?), [gr-qc/0003024](#).
- [34] P. J. E. Peebles, The void phenomenon, *Astrophys. J.* **557** (2001), 495–504.
- [35] S. Gillessen, et al., Monitoring stellar orbits around the massive black hole in the galactic center, *Astrophys. J.* **692** (2009), 1075–1109.
- [36] H. G. Ellis, Cosmic acceleration, inflation, dark matter, and ‘dark energy’ in one neat package, [gr-qc/0701012v3](#).